

# Galilean Creation of the Inflationary Universe

**MASAHIDE YAMAGUCHI**

(Tokyo Institute of Technology)

08/11/15@Hot Topics in General Relativity and Gravitation

arXiv:1504.05710, JCAP 1507 (2015) 07, 017

T. Kobayashi, MY, J. Yokoyama

$$c = \hbar = 1, \quad M_G = 1/\sqrt{8\pi G} \sim 2.4 \times 10^{18} \text{GeV}.$$

# Contents

- **Introduction**

  - Before inflation

  - Violation of null energy condition

  - Galilean Genesis

- **From genesis to inflation (followed by reheating)**

  - Setup (Beyond Horndeski theory)

  - Stability (Powerspectrum of primordial perturbations)

  - Example

- **Summary**

# Introduction


# Inflation

**Inflation**, characterized as **quasi De Sitter expansion**, can naturally solve the problems of the standard big bang cosmology.


- **The horizon problem**
- **The flatness problem**
- **The origin of density fluctuations**
- **The monopole problem**
- ...

# Generic predictions of inflation

- **Spatially flat universe**





- **Almost scale invariant, adiabatic, and Gaussian primordial density fluctuations**
- **Almost scale invariant and Gaussian primordial tensor fluctuations**



**Generates anisotropy of CMBR.**

# Observations of CMB anisotropies

## Planck TT correlation :




Green line : prediction by inflation  
Red points : observation by PLANCK

Angle  $\theta \sim 180^\circ / \ell$

Total energy density  $\leftrightarrow$  Geometry of our Universe

Our Universe is spatially flat !!

## WMAP TE correlation :



Causal seed models

Superhorizon models  
(adiabatic perturbations)

Unfortunately, **primordial tensor perturbations** have not yet been observed.

**What happened before inflation ?**

**and/or**

**How did the Universe begin ?**

# Look back to the past of the Universe

It is often claimed that, if cosmic time goes back to the past, the energy density gets larger and larger, and it eventually reaches the Planck density.

So, unless one completes quantum gravity theory, one cannot discuss the state at the extremely early stage (or even at the onset) of the Universe.

For a perfect fluid :  $T_{\mu\nu} = (\rho + p) u_\mu u_\nu - g_{\mu\nu} p$

The homogeneous and isotropic (Friedmann) Universe:

$$ds^2 = -dt^2 + a^2(t) \gamma_{ij} dx^i dx^j$$

$$\longrightarrow \dot{\rho} = -3H(\rho + p).$$

As long as  $\rho + p \geq 0$  (and  $H > 0$  for the expanding Universe)

$$\longrightarrow \dot{\rho} \leq 0.$$

We do not consider bouncing (contracting) Universe.




# Null energy condition (NEC)

$$T_{\mu\nu}\xi^\mu\xi^\nu \geq 0 \quad \text{for any null vector } \xi^\mu. \\ (g_{\mu\nu}\xi^\mu\xi^\nu = 0)$$

This is the **weakest** among all of the local classical energy conditions.

For a perfect fluid :  $T_{\mu\nu} = (\rho + p)u_\mu u_\nu - g_{\mu\nu}p$



$$\text{NEC} \Leftrightarrow \rho + p \geq 0$$

As long as **the NEC is conserved**, the Universe cannot start from a **low** energy state in the **expanding** Universe.

# How robust is the NEC ?

- Canonical kinetic term with potential:

$$\mathcal{L} = -\frac{1}{2}g^{\mu\nu}\partial_\mu\phi\partial_\nu\phi - V(\phi).$$

$$\longrightarrow \begin{cases} \rho = \frac{1}{2}\dot{\phi}^2 + V(\phi) \\ p = \frac{1}{2}\dot{\phi}^2 - V(\phi) \end{cases} \longrightarrow \rho + p = \dot{\phi}^2 \geq 0.$$

(NEC is conserved)

- How about k-inflation ?

$$\mathcal{L} = K(\phi, X), \quad X = -\frac{1}{2}g^{\mu\nu}\partial_\mu\phi\partial_\nu\phi.$$

$$\longrightarrow \begin{cases} \rho = 2XK_X - X \\ p = K \end{cases} \longrightarrow \rho + p = 2XK_X.$$

$(K_X \equiv \partial K / \partial X)$

Apparently, it looks that, if  $K_X < 0$ , it can violate the NEC.

But, this is not the case.

# Primordial density fluctuations

Garriga & Mukhanov 1999

**Perturbed metric :**

$$ds^2 = -(1 + 2\alpha)dt^2 + 2a^2\partial_i\beta dt dx^i + a^2 e^{2\zeta} dx^2$$

**Comoving gauge :**

$$\phi = \phi(t), \quad \delta\phi = 0.$$

**Prescription:**

- Expand the action up to the second order
- Eliminate  $\alpha$  and  $\beta$  by use of the constraint equations
- Obtain quadratic action for  $\zeta$

➔ 
$$S_S^{(2)} = \int dt d^3x a^3 M_G^2 \frac{\epsilon}{c_s^2} \left( \dot{\zeta}^2 - \frac{c_s^2}{a^2} \zeta_{,k} \zeta_{,k} \right)$$

$$\epsilon = -\frac{\dot{H}}{H^2} = \frac{XK_X}{M_{\text{pl}}^2 H^2}, \quad c_s^2 = \frac{K_X}{K_X + 2XK_{XX}} \quad \text{(sound velocities of curvature perturbations)}$$

In order to avoid the **ghost and gradient instabilities**,  $\epsilon > 0$  &  $c_s^2 > 0$ .

➔ 
$$\rho + p = 2XK_X > 0.$$

(Hsu et al. 2004)  
(See also Dubovsky et al. 2006)

# Stable violation of the NEC

It is impossible to **break the NEC stably** within k-inflation.

$$\left( \mathcal{L} = K(\phi, X), \quad X = -\frac{1}{2}g^{\mu\nu}\partial_\mu\phi\partial_\nu\phi. \right)$$



One may wonder how about introducing **higher derivative terms**.

**Ostrogradski's theorem :**

Assume that  $L = L(q, \dot{q}, \ddot{q})$  and  $\frac{\partial L}{\partial \ddot{q}}$  depends on  $\ddot{q}$  : **(Non-degeneracy)**

$$\frac{\partial L}{\partial q} - \frac{d}{dt} \left( \frac{\partial L}{\partial \dot{q}} \right) + \frac{d^2}{d^2t} \left( \frac{\partial L}{\partial \ddot{q}} \right) = 0, \quad \implies \quad q^{(4)} = q^{(4)}(q^{(3)}, \ddot{q}, \dot{q}, q).$$

This system always leads to **ghost instabilities**.

$$\frac{\partial L}{\partial \phi} - \partial_\mu \left( \frac{\partial L}{\partial (\partial_\mu \phi)} \right) + \partial_\mu \partial_\nu \left( \frac{\partial L}{\partial (\partial_\mu \partial_\nu \phi)} \right) = 0. \quad \implies \quad \frac{i}{(p^2 + m_1^2)(p^2 + m_2^2)} = \frac{1}{m_2^2 - m_1^2} \left( \frac{i}{p^2 + m_1^2} - \frac{i}{p^2 + m_2^2} \right).$$

(propagators)

One loophole to introduce higher derivative terms is that equations of motion should be at most **second order** derivative ones.

# Galileon

Nicolis et al. 2009  
Deffayet et al. 2009

The theory has **Galilean shift symmetry in flat space** :

$$\phi \longrightarrow \phi + c + b_\mu x^\mu \quad (\partial_\mu \phi \longrightarrow \partial_\mu \phi + b_\mu)$$

$$\left\{ \begin{array}{l} \mathcal{L}_1 = \phi \\ \mathcal{L}_2 = (\partial\phi)^2 \\ \mathcal{L}_3 = (\partial\phi)^2 \square\phi \\ \mathcal{L}_4 = (\partial\phi)^2 [(\square\phi)^2 - (\partial_\mu\partial_\nu\phi)^2] \\ \mathcal{L}_5 = (\partial\phi)^2 [(\square\phi)^3 - 3(\square\phi)(\partial_\mu\partial_\nu\phi)^2 + 2(\partial_\mu\partial_\nu\phi)^3] \end{array} \right. \quad \begin{array}{l} (\partial_\mu\partial_\nu\phi)^2 = \partial_\mu\partial_\nu\phi\partial^\mu\partial^\nu\phi, \\ (\partial_\mu\partial_\nu\phi)^3 = \partial_\mu\partial_\nu\phi\partial^\nu\partial^\lambda\phi\partial_\lambda\partial^\mu\phi \end{array}$$

**Lagrangian has higher order derivatives, but EOM is second order.**

Is it possible to **violate the NEC stably** if one includes **higher derivative** terms ?

# Galilean Genesis

Creminelli et al. 2010  
Nicolis et al. 2009

$$S = \int d^4x \sqrt{-g} \left[ \frac{1}{2} M_G^2 R + f^2 e^{2\phi} (\partial\phi)^2 + \frac{f^3}{\Lambda^3} (\partial\phi)^2 \square\phi + \frac{f^3}{2\Lambda^3} (\partial\phi)^4 \right]$$

(In the flat spacetime limit, this theory has **conformal symmetry SO(4,2)**.)


## ● Energy-momentum tensor :

$$\begin{cases} \rho = -f^2 \left( e^{2\phi} \dot{\phi}^2 - \frac{3}{2} \frac{f}{\Lambda^3} \dot{\phi}^4 - 6H \frac{f}{\Lambda^3} \dot{\phi}^3 \right), \\ p = -f^2 \left( e^{2\phi} \dot{\phi}^2 - \frac{1}{2} \frac{f}{\Lambda^3} \dot{\phi}^4 + 2 \frac{f}{\Lambda^3} \dot{\phi}^2 \ddot{\phi} \right). \end{cases}$$

## ● A background solution, (t : $-\infty \rightarrow 0$ ) : **Starts from Minkowski in infinite past.**

$$e^\phi \simeq \frac{1}{\sqrt{2Y_0}} \frac{1}{(-t)}, \quad H \simeq \frac{h_0}{(-t)^3}, \quad \left( a(t) \simeq 1 + \frac{h_0}{2(-t)^2} \right).$$

$$\left( Y_0 \equiv \frac{\Lambda^3}{3f}, \quad h_0 \equiv \frac{1}{2M_G^2} \frac{f^3}{\Lambda^3} \right)$$

  $\rho + p \simeq -\frac{f^3}{\Lambda^3} \frac{4}{(-t)^4} < 0.$  (Actually, you can verify that **H increases**.)

( **The NEC is violated !!** )


# Primordial density fluctuations

**Perturbed metric :**

$$ds^2 = -(1 + 2\alpha)dt^2 + 2a^2\partial_i\beta dt dx^i + a^2 e^{2\zeta} dx^2$$

**Comoving gauge :**

$$\phi = \phi(t), \quad \delta\phi = 0.$$


$$S_S^{(2)} = \int dt d^3x a^3 \left( \mathcal{G}_s \dot{\zeta}^2 - \frac{\mathcal{F}_s}{a^2} \zeta_{,k} \zeta_{,k} \right)$$

In order to avoid the **ghost and gradient instabilities**,  $\mathcal{G}_s > 0$  &  $\mathcal{F}_s > 0$ .

$$\mathcal{G}_s = \mathcal{F}_s \simeq 6M_G^4 \frac{\lambda^3}{f^3} (-t)^2 > 0.$$


( **The NEC is violated stably !!** )

- N.B.**
- A **spectator field like curvaton is responsible for primordial density perturbations** because the genesis field predicts **too blue** ( $n_s \sim 3$ ) perturbations in this simple model.
  - **Primordial tensor perturbations are not generated at first order.**



# Galilean Genesis II

Creminelli et al. 2010  
Nicolis et al. 2009



(figure taken from Creminelli et al. 1007.0027)

- In this scenario, the effective theory **breaks around  $t \sim t_0 = 0$** . So, it is assumed that **the energy density of the genesis field is converted to radiation**, in which hot Universe starts.
- Of course, this is not necessarily a fault of this scenario. A more fundamental theory will be able to describe the transition adequately.


(See 1401.4024 written by Rubakov for good review)



# **From Genesis to inflation**

# From Genesis to inflation

Pirtskhalava et al. 2014



- As a epoch **before inflation** (and the onset of the Universe), use of **Galilean Genesis** is proposed by Pirtskhalava et al.
- Unfortunately, in their concrete construction, **the gradient instabilities appear** during the transition from Genesis to inflation. They are dangerous for large  $k$  modes even during short period because of  $\propto e^{\text{Im}(c_s)kt}$ .

**We try to construct a concrete workable model, in which the Universe starts from Minkowski spacetime in the infinite past, and is smoothly connected to inflation, followed by reheating (graceful exit).**

# Horndeski theory

Horndeski 1974  
Deffayet et al. 2011  
Kobayashi et al. 2011

**Our concrete construction** to realize such a scenario in a healthy way is based on the recent development, **“beyond Horndeski theory”**.

Horndeski theory (= Generalized Galileon) :

$$\left\{ \begin{array}{l} \mathcal{L}_2 = K(\phi, X), \\ \mathcal{L}_3 = -G_3(\phi, X)\square\phi, \\ \mathcal{L}_4 = G_4(\phi, X)R + G_{4X} [(\square\phi)^2 - (\nabla_\mu\nabla_\nu\phi)^2], \\ \mathcal{L}_5 = G_5(\phi, X)G_{\mu\nu}\nabla^\mu\nabla^\nu\phi \\ \quad - \frac{1}{6}G_{5X} [(\square\phi)^3 - 3(\square\phi)(\nabla_\mu\nabla_\nu\phi)^2 + 2(\nabla_\mu\nabla_\nu\phi)^3]. \end{array} \right. \quad X = -\frac{1}{2}(\nabla\phi)^2, \quad G_{iX} \equiv \partial G_i/\partial X.$$

This is the most general (single) scalar-tensor theory which yields **second-order (scalar and gravitational) equations of motion**.

But, in order to avoid the Ostrogradski instabilities, this requirement can be **too strong**. For this purpose, only **time derivatives should be second order** while **spacial ones can be higher**.

# Galileon

Nicolis et al. 2009  
Deffayet et al. 2009

The theory has **Galilean shift symmetry in flat space** :

$$\phi \longrightarrow \phi + c + b_\mu x^\mu \quad (\partial_\mu \phi \longrightarrow \partial_\mu \phi + b_\mu)$$

$$\left\{ \begin{array}{l} \mathcal{L}_1 = \phi \\ \mathcal{L}_2 = (\partial\phi)^2 \\ \mathcal{L}_3 = (\partial\phi)^2 \square\phi \\ \mathcal{L}_4 = (\partial\phi)^2 [(\square\phi)^2 - (\partial_\mu\partial_\nu\phi)^2] \\ \mathcal{L}_5 = (\partial\phi)^2 [(\square\phi)^3 - 3(\square\phi)(\partial_\mu\partial_\nu\phi)^2 + 2(\partial_\mu\partial_\nu\phi)^3] \end{array} \right. \quad \begin{array}{l} (\partial_\mu\partial_\nu\phi)^2 = \partial_\mu\partial_\nu\phi\partial^\mu\partial^\nu\phi, \\ (\partial_\mu\partial_\nu\phi)^3 = \partial_\mu\partial_\nu\phi\partial^\nu\partial^\lambda\phi\partial_\lambda\partial^\mu\phi \end{array}$$


**Lagrangian has higher order derivatives, but EOM is second order.**

**Is it possible to violate the NEC stably if one includes higher derivative terms ?**

# Beyond Horndeski theory

Gleyzes et al. 2014  
Gao 2014

**ADM decomposition:**  $ds^2 = -N^2 dt^2 + \gamma_{ij} (dx^i + N^i dt) (dx^j + N^j dt)$   
( $\varphi = \text{const}$  surfaces)

  $\phi = \phi(t), X = \dot{\phi}^2(t)/(2N^2)$  ( $\varphi$  and  $X$  are functions of  $t$  and  $N$ , and vice versa.)

**Horndeski theory (= Generalized Galileon) :**

$$\mathcal{L} = \sqrt{\gamma} N \sum_a L_a,$$

$$\left\{ \begin{array}{l} L_2 = A_2(t, N), \\ L_3 = A_3(t, N)K, \\ L_4 = A_4(t, N) (K^2 - K_{ij}^2) + B_4(t, N)R^{(3)}, \\ L_5 = A_5(t, N) (K^3 - 3KK_{ij}^2 + 2K_{ij}^3) + B_5(t, N)K^{ij} \left( R_{ij}^{(3)} - \frac{1}{2}g_{ij}R^{(3)} \right). \end{array} \right.$$

**with**  $A_4 = -B_4 - N \frac{\partial B_4}{\partial N}, \quad A_5 = \frac{N}{6} \frac{\partial B_5}{\partial N}.$

$$\left\{ \begin{array}{l} \mathcal{L}_2 = K(\phi, X), \\ \mathcal{L}_3 = -G_3(\phi, X)\square\phi, \\ \mathcal{L}_4 = G_4(\phi, X)R + G_{4X} [(\square\phi)^2 - (\nabla_\mu \nabla_\nu \phi)^2], \\ \mathcal{L}_5 = G_5(\phi, X)G_{\mu\nu} \nabla^\mu \nabla^\nu \phi \\ \quad - \frac{1}{6}G_{5X} [(\square\phi)^3 - 3(\square\phi)(\nabla_\mu \nabla_\nu \phi)^2 + 2(\nabla_\mu \nabla_\nu \phi)^3]. \end{array} \right.$$

**Kij : extrinsic curvature**  
**Rij(3) : intrinsic curvature**

**Gleyzes et al. (GLPV) pointed out that, even if the above two relations are absent, the number of the propagating degrees of freedom remains unchanged. Gao showed that further extension is possible.**

# Our setup

$$\mathcal{L} = \sqrt{\gamma} N \sum_a L_a$$

$$\left\{ \begin{array}{l} L_2 = A_2(t, N), \\ L_3 = A_3(t, N)K, \\ L_4 = A_4(t, N) \left( \lambda_1 K^2 - K_{ij}^2 \right) + B_4(t, N)R^{(3)}, \\ L_5 = A_5(t, N) \left( \lambda_2 K^3 - 3\lambda_3 K K_{ij}^2 + 2K_{ij}^3 \right) \\ \quad + B_5(t, N)K^{ij} \left( R_{ij}^{(3)} - \frac{1}{2}g_{ij}R^{(3)} \right). \end{array} \right.$$

( The **GLPV theory** corresponds to the case with  $\lambda_1 = \lambda_2 = \lambda_3 = 1$ .)

$$ds^2 = -N^2 dt^2 + \gamma_{ij} (dx^i + N^i dt) (dx^j + N^j dt)$$

$$\left\{ \begin{array}{l} N = \bar{N}(t) (1 + \delta n), \\ N_i = \bar{N}(t) \partial_i \chi, \\ \gamma_{ij} = a^2(t) e^{2\zeta} (e^h)_{ij}. \end{array} \right.$$

curvature perturbations

tensor perturbations

( $h_{ii} = h_{ij,j} = 0$ )

# Perturbations

● **Tensor perturbations :**

$$\mathcal{L}_T^{(2)} = \frac{\bar{N}a^3}{8} \left[ \frac{\mathcal{G}_T}{\bar{N}^2} \dot{h}_{ij}^2 - \frac{\mathcal{F}_T}{a^2} (\partial h_{ij})^2 \right] \quad \begin{cases} \mathcal{G}_T := -2A_4 - 6(3\lambda_3 - 2)A_5H, \\ \mathcal{F}_T := 2B_4 + \frac{1}{\bar{N}} \frac{dB_5}{dt}. \end{cases}$$

$$\left( H := \frac{\dot{a}}{(\bar{N}a)} \right)$$

● **Curvature perturbations : (spatial higher derivative appears !!)**

$$\mathcal{L}_S^{(2)} = \bar{N}a^3 \left[ \mathcal{G}_S \frac{\zeta^2}{\bar{N}^2} + \zeta \left( \mathcal{F}_S \frac{\partial^2}{a^2} - \mathcal{H}_S \frac{\partial^4}{a^4} \right) \zeta \right] \quad \longrightarrow \quad \omega^2 = \frac{\mathcal{F}_S}{\mathcal{G}_S} k^2 + \frac{\mathcal{H}_S k^4}{\mathcal{G}_S a^2}.$$

$$\left\{ \begin{array}{l} \mathcal{G}_S := \frac{\Sigma \mathcal{G}_T^2}{\Theta^2 + \Sigma \mathcal{C}} + 3\mathcal{G}_T, \\ \mathcal{F}_S := \frac{1}{\bar{N}a} \frac{d}{dt} \left( \frac{a\Theta \mathcal{G}_B \mathcal{G}_T}{\Theta^2 + \Sigma \mathcal{C}} \right) - \mathcal{F}_T, \\ \mathcal{H}_S := \frac{\mathcal{G}_B^2 \mathcal{C}}{\Theta^2 + \Sigma \mathcal{C}}. \end{array} \right. \quad \left\{ \begin{array}{l} \Sigma := \bar{N}A_2' + \frac{1}{2}\bar{N}^2 A_2'' + \frac{3}{2}\bar{N}^2 A_3'' H \\ \quad + 3\eta_4 (2A_4 - 2\bar{N}A_4' + \bar{N}^2 A_4'') H^2 \\ \quad + 3\eta_5 (6A_5 - 4\bar{N}A_5' + \bar{N}^2 A_5'') H^3, \\ \Theta := \frac{\bar{N}A_3'}{2} - 2\eta_4 (A_4 - \bar{N}A_4') H \\ \quad - 3\eta_5 (2A_5 - \bar{N}A_5') H^2, \\ \mathcal{G}_A := -2\eta_4 A_4 - 6\eta_5 A_5 H, \\ \mathcal{G}_B := 2(B_4 + \bar{N}B_4') - H\bar{N}B_5', \\ \mathcal{C} := (1 - \lambda_1)A_4 - (6 + 9\lambda_2 - 15\lambda_3)A_5 H. \end{array} \right.$$

$$(\eta_4 := (3\lambda_1 - 1)/2, \quad \eta_5 := (9\lambda_2 - 9\lambda_3 + 2)/2)$$

**N.B.** ● **C = 0** for  $\lambda_1 = \lambda_2 = \lambda_3 = 1$ .

● **Even if  $F_s < 0$  (with  $G_s > 0$ ), the curvature perturbations with large  $k$  are stabilized for  $H_s > 0$ .**



# Concrete example

$$\left\{ \begin{array}{l} A_2 = M_2^4 f^{-2(\alpha+1)}(t) a_2(N), \\ A_3 = M_3^3 f^{-(2\alpha+1)}(t) a_3(N), \\ A_4 = -\frac{M_G^2}{2} + M_4^2 f^{-2\alpha}(t) a_4(N), \\ A_5 = M_5 f(t) a_5(N), \end{array} \right. \quad (\alpha > 0) \quad (\varphi, \mathbf{X}) \rightarrow (t, \mathbf{N})$$

$$\left\{ \begin{array}{l} L_2 = A_2(t, N), \\ L_3 = A_3(t, N)K, \\ L_4 = A_4(t, N) (\lambda_1 K^2 - K_{ij}^2) + B_4(t, N)R^{(3)}, \\ L_5 = A_5(t, N) (\lambda_2 K^3 - 3\lambda_3 K K_{ij}^2 + 2K_{ij}^3) \\ \quad + B_5(t, N)K^{ij} \left( R_{ij}^{(3)} - \frac{1}{2}g_{ij}R^{(3)} \right). \end{array} \right.$$

**Background dynamics :**  $\mathcal{L}^{(0)} = \bar{N} a^3 (A_2 + 3A_3 H + 6\eta_4 A_4 H^2 + 6\eta_5 A_5 H^3)$ .

$$\left\{ \begin{array}{l} -\mathcal{E} := (\bar{N} A_2)' + 3\bar{N} A_3' H + 6\eta_4 \bar{N}^2 (\bar{N}^{-1} A_4)' H^2 + 6\eta_5 \bar{N}^3 (\bar{N}^{-2} A_5)' H^3 = 0, \\ \mathcal{P} := A_2 - 6\eta_4 A_4 H^2 - 12\eta_5 A_5 H^3 - \frac{1}{\bar{N}} \frac{d}{dt} (A_3 + 4\eta_4 A_4 H + 6\eta_5 A_5 H^2) = 0. \end{array} \right.$$

( $' := d/d\bar{N}$ )

• **Genesis phase ( $t < t_0$ ) :**  $f(t) \simeq \dot{f}_0 t$  ( $\dot{f}_0 = \text{const} < 0$ )

$$\rightarrow \left\{ \begin{array}{l} \bar{N} \simeq N_0 (= \text{const}) \text{ with } a_2(N_0) + N_0 a_2'(N_0) = 0. \\ H = -\frac{\hat{p}}{2(2\alpha+1)\eta_4 M_G^2 |\dot{f}_0|} \frac{N_0}{f_0} f^{-(2\alpha+1)} \sim \frac{1}{(-t)^{2\alpha+1}}, \\ a = 1 - \frac{\hat{p}}{4\alpha(2\alpha+1)\eta_4 M_G^2} \frac{N_0^2}{\dot{f}_0^2} f^{-2\alpha}. \end{array} \right. \quad \left( \hat{p} = M_2^4 a_2(N_0) + (2\alpha+1) M_3^3 a_3(N_0) \frac{\dot{f}_0}{N_0} \right)$$

(The background dynamics for  $\alpha = 1$  coincides with that of the original Genesis model.)

# Concrete example II

- Inflationary phase (  $t_{end} > t > t_0$  ) :  $f(t) \simeq f_1 (= \text{const})$

→ 
$$\begin{cases} \bar{N} \simeq N_{\text{inf}} (= \text{const}), \\ H \simeq H_{\text{inf}} (= \text{const}). \end{cases}$$

with 
$$\begin{cases} -\mathcal{E} = (N_{\text{inf}} A_2)' + 3N_{\text{inf}} A_3' H_{\text{inf}} + 6\eta_4 N_{\text{inf}}^2 (N_{\text{inf}}^{-1} A_4)' H_{\text{inf}}^2 \\ \quad + 6\eta_5 N_{\text{inf}}^3 (N_{\text{inf}}^{-2} A_5)' H_{\text{inf}}^3 = 0, \\ \mathcal{P} = A_2 - 6\eta_4 A_4 H_{\text{inf}}^2 - 12\eta_5 A_5 H_{\text{inf}}^3 = 0. \end{cases}$$

**N.B.** A weak time dependence of  $f(t)$  yields slight deviation from exact DS.

- Graceful exit (  $t > t_{end}$  ) :  $f(t) \sim t^{1/(\alpha+1)}$

**If  $m > 4$ , gravitational reheating can work.**

→ 
$$\begin{cases} \bar{N} \simeq N_e (= \text{const}), \\ H^2 \sim 1/t^2 \sim f^{-2(\alpha+1)} \propto 1/a^m \quad (m := 3N_e a_2' / (N_e a_2)' > 0). \end{cases}$$

with 
$$\begin{cases} -\mathcal{E} = (N_e A_2)' + 3\eta_4 M_G^2 H^2 + \mathcal{O}(f^{-(3\alpha+2)}) = 0, \\ \mathcal{P} = A_2 + 3\eta_4 M_G^2 H^2 + \frac{2\eta_4 M_G^2}{N_e} \frac{dH}{dt} + \mathcal{O}(f^{-(3\alpha+2)}) = 0. \end{cases}$$

# Perturbations

● **Tensor perturbations :**

$$\mathcal{L}_T^{(2)} = \frac{\bar{N}a^3}{8} \left[ \frac{\mathcal{G}_T}{\bar{N}^2} \dot{h}_{ij}^2 - \frac{\mathcal{F}_T}{a^2} (\partial h_{ij})^2 \right] \quad \begin{cases} \mathcal{G}_T := -2A_4 - 6(3\lambda_3 - 2)A_5H, \\ \mathcal{F}_T := 2B_4 + \frac{1}{\bar{N}} \frac{dB_5}{dt}. \end{cases}$$

$$\left( H := \frac{\dot{a}}{(\bar{N}a)} \right)$$

● **Curvature perturbations : (spatial higher derivative appears !!)**

$$\mathcal{L}_S^{(2)} = \bar{N}a^3 \left[ \mathcal{G}_S \frac{\zeta^2}{\bar{N}^2} + \zeta \left( \mathcal{F}_S \frac{\partial^2}{a^2} - \mathcal{H}_S \frac{\partial^4}{a^4} \right) \zeta \right] \quad \longrightarrow \quad \omega^2 = \frac{\mathcal{F}_S}{\mathcal{G}_S} k^2 + \frac{\mathcal{H}_S k^4}{\mathcal{G}_S a^2}.$$

$$\left\{ \begin{array}{l} \mathcal{G}_S := \frac{\Sigma \mathcal{G}_T^2}{\Theta^2 + \Sigma \mathcal{C}} + 3\mathcal{G}_T, \\ \mathcal{F}_S := \frac{1}{\bar{N}a} \frac{d}{dt} \left( \frac{a\Theta \mathcal{G}_B \mathcal{G}_T}{\Theta^2 + \Sigma \mathcal{C}} \right) - \mathcal{F}_T, \\ \mathcal{H}_S := \frac{\mathcal{G}_B^2 \mathcal{C}}{\Theta^2 + \Sigma \mathcal{C}}. \end{array} \right. \quad \left\{ \begin{array}{l} \Sigma := \bar{N}A_2' + \frac{1}{2}\bar{N}^2 A_2'' + \frac{3}{2}\bar{N}^2 A_3'' H \\ \quad + 3\eta_4 (2A_4 - 2\bar{N}A_4' + \bar{N}^2 A_4'') H^2 \\ \quad + 3\eta_5 (6A_5 - 4\bar{N}A_5' + \bar{N}^2 A_5'') H^3, \\ \Theta := \frac{\bar{N}A_3'}{2} - 2\eta_4 (A_4 - \bar{N}A_4') H \\ \quad - 3\eta_5 (2A_5 - \bar{N}A_5') H^2, \\ \mathcal{G}_A := -2\eta_4 A_4 - 6\eta_5 A_5 H, \\ \mathcal{G}_B := 2(B_4 + \bar{N}B_4') - H\bar{N}B_5', \\ \mathcal{C} := (1 - \lambda_1)A_4 - (6 + 9\lambda_2 - 15\lambda_3)A_5 H. \end{array} \right.$$

$$(\eta_4 := (3\lambda_1 - 1)/2, \quad \eta_5 := (9\lambda_2 - 9\lambda_3 + 2)/2)$$

**N.B.** ● **C = 0** for  $\lambda_1 = \lambda_2 = \lambda_3 = 1$ .




● **Even if  $F_s < 0$  (with  $G_s > 0$ ), the curvature perturbations with large  $k$  are stabilized for  $H_s > 0$ .**

# Numerical calculations

From Genesis to inflation :

$$f(t) = \frac{f_0}{2} \left\{ t - \frac{\ln[2 \cosh(st)]}{s} \right\} + f_1,$$

$$f_0 = -10^{-1}, \quad f_1 = 10, \quad s = 2 \times 10^{-3} \simeq t_0^{-1}$$



During short period,  
 $\mathcal{F}_S$  becomes negative.  
 But, the perturbations  
 for large  $k$  are stabilized  
 thanks to the  $k^4$  terms.

The perturbations for  
 small  $k$  grow during  
 short period, but growth  
 is mild and finite.

The situation is similar  
 to the transition from  
 inflation to RD.

FIG. 2: The background evolution of the Hubble parameter  $H$  (a) and the lapse function  $N$  (b) around the genesis-de Sitter transition.

FIG. 3: The sound speed squared  $\mathcal{F}_S/\mathcal{G}_S$  (a) and the coefficient of  $k^4$  (divided by  $\mathcal{G}_S$ ) (b) around the genesis-de Sitter transition.

# Conclusions

- We constructed a concrete example **from Galilean Genesis to inflationary phase followed by graceful exit**, based on the recent development **beyond the Horndeski theory**.
- **The sound velocities squared (or  $F_s$ )** during transitions from Genesis to inflation and from inflation to RD **become negative** for a short period.
- But thanks to a **non-trivial dispersion relation** coming from the fourth order derivative term in the quadratic action, **modes with higher  $k$  are completely stable and the growth of perturbations with smaller  $k$  is finite and controllable**.
- Our model can describe a Genesis scenario with graceful exit **(even without inflationary phase)**, in which no (first order) primordial tensor perturbations are produced. The **detection or non-detection** of primordial tensor perturbations may discriminate Genesis scenarios with or without inflation.