

EXACT BLACK HOLE SOLUTIONS IN MASSIVE GRAVITY

M.Bebronne¹, P. Tinyakov^{1,2}

¹ Université Libre de Bruxelles (ULB)
Brussels

² Institute for Nuclear Research (INR)
Moscow

[arXiv:0902.3899]

SW3

April 2009, Cargese

Motivation

The model

Spherical
equations

Analytical
solution

Summary

Outline

Motivation

The model

Spherical equations

Analytical solution

Summary

**EXACT BLACK
HOLE
SOLUTIONS IN
MASSIVE
GRAVITY**

M.Bebroune,
P. Tinyakov

Motivation

The model

**Spherical
equations**

**Analytical
solution**

Summary

Why study modifications of gravity?

**EXACT BLACK
HOLE
SOLUTIONS IN
MASSIVE
GRAVITY**

M.Bebbronne,
P. Tinyakov

- ▶ Gravity is the most mysterious force which is quite poorly understood
 - ▶ GR is a beautiful and unique classical theory. However:
 - ▶ problems at quantum level
 - ▶ problems at largest scales
- ⇒ better understanding of gravity is needed
- ▶ Such an understanding may come from attempts to construct a consistent modified theory of gravity
 - ▶ To be of phenomenological interest, the model has to coincide with GR at scales from mm to kpc

Motivation

The model

**Spherical
equations**

**Analytical
solution**

Summary

Why black holes are of particular interest in such models

EXACT BLACK HOLE SOLUTIONS IN MASSIVE GRAVITY

M.Bebbronne,
P. Tinyakov

- ▶ Black holes are the most fascinating non-linear manifestations of gravity. Hence they are good theoretical laboratories for studying gravity models.
- ▶ The properties of black holes are deeply interrelated with fundamental structure of the underlying theory.
- ▶ Black holes are states where macroscopic and microscopic features of theory (including quantum properties) are mixed together.
- ▶ Finally, BH's most likely exist in nature and will be observed and perhaps studied at the quantitative level in the near future.

Motivation

The model

Spherical
equations

Analytical
solution

Summary

Massive gravity as a “deformation” of GR

EXACT BLACK
HOLE
SOLUTIONS IN
MASSIVE
GRAVITY

M.Bebbronne,
P. Tinyakov

- ▶ Massive gravity: scalar-tensor theory with Lorentz invariance that is spontaneously broken by space-time dependent condensates of scalar fields. The action is [Dubovsky'2004]

$$S = \int \sqrt{g} \left\{ -R + L(\text{matter}) + \Lambda^4 F(\text{scalars}) \right\}$$

Note: no direct coupling between scalars and ordinary matter

- ▶ The model is understood as an effective theory below the scale Λ

Motivation

The model

Spherical
equations

Analytical
solution

Summary

Coupling of scalars to gravity

- Four scalars ϕ^0, ϕ^i with derivative coupling

$$F = F(X, W^{ij})$$

$$X = g^{\mu\nu} \partial_\mu \phi^0 \partial_\nu \phi^0$$

$$W^{ij} = g^{\mu\nu} \partial_\mu \phi^i \partial_\nu \phi^j - g^{\mu\nu} \partial_\mu \phi^0 \partial_\nu \phi^i \cdot g^{\alpha\beta} \partial_\alpha \phi^0 \partial_\beta \phi^j / X$$

- Obeys the symmetry

$$\phi^i \rightarrow \phi^i + \xi^i(\phi^0)$$

which ensures the absence of ghosts and instabilities at the perturbative level.

Motivation

The model

Spherical
equations

Analytical
solution

Summary

Vacuum and perturbations

- ▶ For a generic function F , the energy-momentum of scalars $T_{\mu\nu}$ vanishes for the following configuration

$$\begin{aligned}g_{\mu\nu} &= \eta_{\mu\nu} \\ \phi^0 &= \Lambda^2 \cdot t \\ \phi^i &= \Lambda^2 \cdot x^i\end{aligned}$$

\implies Minkowski space is a solution

- ▶ The spectrum of perturbations contains massive tensor modes and no other particles. Graviton mass is:

$$m \sim \Lambda^2 / M_{\text{Pl}}$$

Motivation

The model

Spherical
equations

Analytical
solution

Summary

Cosmological solutions

EXACT BLACK HOLE SOLUTIONS IN MASSIVE GRAVITY

M. Bebronne,
P. Tinyakov

- Flat cosmological ansatz:

$$ds^2 = dt^2 - a^2(t) dx_i^2$$

$$\phi^0 = \Lambda^2 \cdot \phi(t)$$

$$\phi^i = \Lambda^2 \cdot x^i$$

- Cosmological evolution has an attractor where $F = F(X^\gamma W^{ij})$. In that point the scalars give two contributions to ρ , behaving like cosmological constant and like matter with the equation of state depending on γ .

Motivation

The model

Spherical
equations

Analytical
solution

Summary

Summary of the model:

- Spectrum of perturbations: massive graviton (2 polarizations) with

$$m \sim \Lambda^2 / M_{\text{Pl}}$$

Current bound $m \lesssim 10^{-20}$ eV [Dubovsky,PT,Tkachev'2005]

- Force between static sources: at the attractor point, standard Newton's law + strongly suppressed corrections
- Cosmology: flat Universe; extra contributions from scalar fields behaving like cosmological constant and matter with the equation of state $-1 < w < \infty$
- Cosmological perturbations: standard behavior in some range of parameters [Bebronne,PT'2007]
- !Rem: to be tested against BBN

Do black holes get modified in this model?

- ▶ Schwarzschild metric is a solution to the field equations with

$$\phi_0 = \Lambda^2 \left(t + 2\sqrt{rr_s} + r_s \ln \frac{\sqrt{r} - \sqrt{r_s}}{\sqrt{r} + \sqrt{r_s}} \right)$$

[Dubovsky,PT,Zaldarriaga'2007]

- ▶ Kerr metric is not a solution no matter what are scalar fields
- ▶ Generically, black holes may have hair due to presence of instantaneous interaction
- ▶ Modified black hole solutions have been found in bi-gravity models *[Berezhiani,Comelli,Nesti,Pilo'2008]*

Motivation

The model

Spherical
equations

Analytical
solution

Summary

Spherically-symmetric ansatz

EXACT BLACK
HOLE
SOLUTIONS IN
MASSIVE
GRAVITY

M. Bebronne,
P. Tinyakov

6 functions to begin with:

$$ds^2 = \alpha(r) dt^2 + 2\delta(r) dt dr - \beta(r) dr^2 - \kappa(r) d\Omega^2$$

$$\phi^0 = \Lambda^2(t + h(r))$$

$$\phi^i = \phi(r) \frac{\Lambda^2 x^i}{r}$$

Use gauge freedom to set

$$\kappa(r) = r^2$$

$$\delta(r) = 0$$

\Rightarrow 4 radial functions $\alpha(r)$, $\beta(r)$, $h(r)$ and $\phi(r)$

Motivation

The model

Spherical
equations

Analytical
solution

Summary

Einstein equations

EXACT BLACK HOLE SOLUTIONS IN MASSIVE GRAVITY

M. Bebronne,
P. Tinyakov

Motivation

The model

Spherical
equations

Analytical
solution

Summary

$$G_0^0 = \frac{1}{M_{Pl}^2} T_0^0 \quad G_r^r = \frac{1}{M_{Pl}^2} T_r^r$$
$$G_\theta^\theta = \frac{1}{M_{Pl}^2} T_\theta^\theta \quad 0 = T_0^r$$

When the last equation holds, $T_0^0 = T_r^r$, which implies that $G_0^0 = G_r^r$, which in turn implies that

$$\alpha(r)\beta(r) = 1.$$

We are left with 3 differential equations for 3 variables.

Analytical example

Main tool — choice of the function F .

From rotational invariance, F has to depend on W^{ij} through three combinations $w_n = \text{Tr}(W^{ij})^n$. Take

$$F = c_0 \left(\frac{1}{X} + w_1 \right) + c_1 \left(w_1^3 - 3w_1 w_2 - 6w_1 + 2w_3 - 12 \right)$$

Constraints to be satisfied:

- ▶ Minkowski space has to be a solution
- ▶ Graviton should be non-tachyonic

$$c_0 - 6c_1 \geq 0$$

- ▶ Stability upon addition of higher-derivative terms

$$c_0 > 0$$

Solution

EXACT BLACK HOLE SOLUTIONS IN MASSIVE GRAVITY

M.Bebbronne,
P. Tinyakov

Due to specific choice of F , equation $T_0^r = 0$ simplifies:

$$\frac{1}{X} \left[-c_0 + \phi'^2 \left(6c_1 \frac{\phi^4}{r^4} + c_0 - 6c_1 \right) \right] = 0$$

The solution to this equation is

$$\phi = br,$$

with

$$(b^2 - 1)(6b^4 + 6b^2 + c_0/c_1) = 0$$

Motivation

The model

Spherical
equations

Analytical
solution

Summary

Solution

The remaining two equations can also be solved analytically,

$$\alpha(r) = 1 - \frac{2M}{r} - \frac{S}{r^\lambda} + \Lambda_c r^2$$

$$\beta(r) = 1/\alpha(r)$$

$$h(r) = \pm \int \frac{dr}{\alpha} \left[1 - \alpha \left(\frac{S}{c_0 m^2} \frac{\lambda - 1}{r^{\lambda+2}} + 1 \right)^{-1} \right]^{1/2}$$

$$\phi(r) = b \cdot r$$

where

$$\lambda = -12b^6 \frac{c_1}{c_0}$$


$$\Lambda_c = 2m^2 c_1 (b^6 - 1)$$

Behavior of solutions

Case of interest: $c_1 = -1$, $c_0 > 0 \implies \lambda > 0$.

Consider the branch $\Lambda_c = 0$ (no cosmological constant)


$M > 0$, $S < 0$.



Gravity is weaker close to horizon

Behavior of solutions

$$M > 0, S < 0.$$



**EXACT BLACK
HOLE
SOLUTIONS IN
MASSIVE
GRAVITY**

M. Bebronne,
P. Tinyakov

Motivation

The model


Spherical
equations

Analytical
solution

Summary

Behavior of solutions

$$M < 0, S > 0.$$



Gravity is stronger close to horizon

**EXACT BLACK
HOLE
SOLUTIONS IN
MASSIVE
GRAVITY**

M. Bebronne,
P. Tinyakov

Motivation

The model

Spherical
equations

Analytical
solution

Summary

Summary

EXACT BLACK HOLE SOLUTIONS IN MASSIVE GRAVITY

M.Bebroune,
P. Tinyakov

Motivation

The model

Spherical
equations

Analytical
solution

Summary

- ▶ Exact spherically-symmetric solutions in modified gravity are different from Schwarzschild black holes
- ▶ They show rich behavior, including anti-gravitation at large distances
- ▶ Open questions:
 - ▶ stability
 - ▶ formation of configurations with $S \neq 0$