

Analytic Infinite Derivative (AID) gravity
or
String Field Theory derivation of non-local cosmology

Alexey Koshelev

Universidade da Beira Interior

Cargese, May 1, 2017

Based on arXiv:1604.01440 in collaboration with S. Korumilli, P. Moniz,
arXiv:1604.03127 in collaboration with L.Modesto, L.Rachwal and A.Starobinsky
and the current work in progress

Motivation

- Gravity is not renormalizable
- Stelle's 1977 and 1978 papers show that R^2 gravity is renormalizable gravity with the price of a physical (Weyl) ghost
(- + ++) $\Rightarrow -\partial\varphi^2$ is a good field
- Ostrogradski statement from 1850 forbids higher derivatives in general
- Starobinsky inflation is based on R^2 and works perfectly

Exorcising ghosts

- Constrained systems like GR
- Special theories like Horndeski models
- Infinitely many derivatives

String Field Theory in 5 min

- Solid stuff
- Unitary
- Finite
- Describes all our fields universally
- From the embedding space-time point of view contains analytic infinite derivative operators
- Well, has its own unsolved puzzles

The most general action to consider

We start with

$$S = \int d^4x \sqrt{-g} \left(\mathcal{P}_0 + \sum_i \mathcal{P}_i \prod_I (\hat{\mathcal{O}}_{iI} \mathcal{Q}_{iI}) \right)$$

Here \mathcal{P} and \mathcal{Q} depend on curvatures and \mathcal{O} are operators made of covariant derivatives.

We are looking for the most general action which captures in full generality the properties of a linearized model around *maximally symmetric space-times* given $g_{\mu\nu} = \bar{g}_{\mu\nu} + h_{\mu\nu}$.

The result is [\[arxiv.1602.08475\]](#)

$$S = \int d^4x \sqrt{-g} \left(\frac{M_P^2 R}{2} + \frac{\lambda}{2} \left(R \mathcal{F}(\square) R + L_{\mu\nu} \mathcal{F}_L(\square) L^{\mu\nu} + W_{\mu\nu\lambda\sigma} \mathcal{F}_W(\square) W^{\mu\nu\lambda\sigma} \right) - \Lambda \right)$$

Here $L_{\mu\nu} = R_{\mu\nu} - \frac{1}{4} R g_{\mu\nu}$.

Quadratic action around (A)dS with $\bar{R} = 4\Lambda/M_P^2$

The covariant decomposition is

$$h_{\mu\nu} = \frac{2}{M_P^2} h_{\mu\nu}^\perp + \bar{\nabla}_\mu A_\nu + \bar{\nabla}_\nu A_\mu + (\bar{\nabla}_\mu \bar{\nabla}_\nu - \frac{1}{4} \frac{2}{M_P^2} \sqrt{\frac{8}{3}} \bar{g}_{\mu\nu} \bar{\square}) B + \frac{1}{4} \frac{2}{M_P^2} \sqrt{\frac{8}{3}} \bar{g}_{\mu\nu} h$$

Here $\bar{\nabla}^\mu h_{\mu\nu}^\perp = \bar{g}^{\mu\nu} h_{\mu\nu}^\perp = \bar{\nabla}^\mu A_\mu = 0$.

Spin-2:

$$S_2 = \frac{1}{2} \int dx^4 \sqrt{-\bar{g}} h_{\nu\mu}^\perp \left(\bar{\square} - \frac{\bar{R}}{6} \right) [\mathcal{P}(\bar{\square})] h^{\perp\mu\nu}$$

$$\mathcal{P}(\bar{\square}) = 1 + \frac{2}{M_P^2} \lambda f_0 \bar{R} + \frac{\lambda}{M_P^2} \left\{ \mathcal{F}_L(\bar{\square}) \left(\bar{\square} - \frac{\bar{R}}{6} \right) + 2\mathcal{F}_W \left(\bar{\square} + \frac{\bar{R}}{3} \right) \left(\bar{\square} - \frac{\bar{R}}{3} \right) \right\}$$

Spin-0 (here $\phi \equiv \bar{\square} B - h$):

$$S_0 = -\frac{1}{2} \int dx^4 \sqrt{-\bar{g}} \phi (3\bar{\square} + \bar{R}) [\mathcal{S}(\bar{\square})] \phi$$

$$\mathcal{S}(\bar{\square}) = 1 + \frac{2}{M_P^2} \lambda f_0 \bar{R} - \frac{\lambda}{M_P^2} \left\{ 2\mathcal{F}(\bar{\square}) (3\bar{\square} + \bar{R}) + \frac{1}{2} \mathcal{F}_L \left(\bar{\square} + \frac{2}{3} \bar{R} \right) \bar{\square} \right\}$$

Physical excitations

Effectively we modify the propagators as follows

$$\square - m^2 \rightarrow \mathcal{G}(\square)$$

To preserve the physics we demand

$$\mathcal{G}(\square) = (\square - m^2)e^{\sigma(\square)}$$

Here $\sigma(\square)$ must be an *entire* function resulting that the exponent of it has no roots.

We arrange this in our model by virtue of functions \mathcal{F} . At this stage we can drop any one of three \mathcal{F} -s. The simplest choice is to drop \mathcal{F}_L .

Starobinsky inflation in non-local gravity

For any:

$$\square R = r_1 R + r_2$$

We have a solution:

$$\mathcal{F}^{(1)}(r_1) = 0, \quad \frac{r_2}{r_1}(\mathcal{F}(r_1) - f_0) = -\frac{M_P^2}{2\lambda} + 3r_1\mathcal{F}(r_1), \quad 4\Lambda r_1 = -r_2 M_P^2$$

In the case of interest $\Lambda = 0$.

Notice that the we have started with the trace of Einstein equations in a local R^2 gravity.

Saying local gravity we do mean *any* including pathological parameters in that local counterpart.

Choice of $\mathcal{F}(\square)$

We should arrange that the theory is ghost-free meaning that no more than one pole arises in the scalar sector. The new degree of freedom is named scalaron and its mass is denoted as M . A possible form is:

$$\frac{\lambda}{M_P^2} \mathcal{F}(\square) = -\frac{1}{6\square} \left[e^{H_0(\square)} \left(1 - \frac{\square}{M_P^2} \right) - 1 \right]$$

The conditions on $\mathcal{F}(\square)$ imply that $H_0(\square)$ is an entire function and moreover:

$$r_1 = M^2$$

$$H_0(r_1) = 0$$

Power spectra and r

Tensor modes

$$|\delta_h|^2 = \frac{H^2}{2\pi^2 \lambda \mathcal{F}_1 \bar{R}} e^{2\omega(\bar{R}/6)} \quad \text{where } \mathcal{P}(\bar{\square}) = e^{2\omega(\bar{\square})}$$

Scalar modes (actually $\mathcal{R} = \Psi + \frac{H}{\dot{R}} \delta R_{GI}$)

$$|\delta_{\mathcal{R}}|^2 \approx \frac{H^6}{16\pi^2 \dot{H}^2 3\lambda \mathcal{F}_1 \bar{R}}$$

Tensor to scalar ratio r

$$r = \frac{2|\delta_h|^2}{|\delta_{\mathcal{R}}|^2} = 48 \frac{\dot{H}^2}{H^4} e^{2\omega(\bar{R}/6)}$$

All quantities here are at the horizon crossing $k = Ha$.

Analogously

$$N = \int_{t_i}^{t_f} H dt = \frac{1}{2\epsilon_1} \Rightarrow r = 48\epsilon_1^2 e^{2\omega(\bar{R}/6)} = \frac{12}{N^2} e^{2\omega(\bar{R}/6)}$$

UV completeness

Minkowski propagator:

$$\Pi = - \left(\frac{P^{(2)}}{k^2 e^{H_2(-k^2)}} - \frac{P^{(0)}}{2k^2 e^{H_0(-k^2)} \left(1 + \frac{k^2}{M^2}\right)} \right)$$

To guarantee that the QFT machinery works we arrange a polynomial decay of the propagator near infinity. The rate of the decay is our choice. Recall that we still need the functions $H_{0,2}$ to be entire.

An example of such a function can be, for instance

$$H \sim \Gamma(0, p(z)^2) + \gamma_E + \log(p(z)^2)$$

where $p(z)$ is a polynomial.

Beyond 1-loop the powercounting arguments work just like in the higher derivative regularization.

p-adic reformulation of the non-local gravity

The scalar part of the previous action is equivalent to the following one

$$S = \int d^4x \sqrt{-g} \left(\frac{M_P^2 R}{2} \left(1 + \frac{2}{M_P^2} \psi \right) - \frac{1}{2\lambda} \psi \frac{1}{\mathcal{F}(\square)} \psi + \dots \right)$$

An important property here is the non-minimal coupling of a scalar field to gravity.

The conformal transform $\left(1 + \frac{2}{M_P^2} \psi \right)^2 g_{\mu\nu} = \bar{g}_{\mu\nu}$ allows us to decouple the gravity and the scalar field even more

$$S = \int d^4x \sqrt{-\bar{g}} \left(\frac{M_P^2 \bar{R}}{2} - \frac{M_P^2}{2} \frac{6}{(M_P^2 + 2\psi)^2} \bar{g}^{\mu\nu} \partial_\mu \psi \partial_\nu \psi - \frac{M_P^4}{2\lambda (M_P^2 + 2\psi)^2} \psi \mathcal{G}(\mathcal{D}) \psi \right)$$

Here

$$\mathcal{G}(\mathcal{D}) = \frac{1}{\mathcal{F}(\mathcal{D})} \text{ and } \mathcal{D} = \left(1 + \frac{2}{M_P^2} \psi \right) \square_{\bar{g}} - \frac{2}{M_P^2} \bar{g}^{\mu\nu} \partial_\mu \psi \partial_\nu$$

Conclusions

- A UV complete and unitary gravity is presented
- Starobinsky inflation is natively embedded in this model
- The theory predicts a modified value for r
- A connection to p -adic strings is outlined
- Few words about SFT in this story, Slavnov-Taylor identities, Cutkosky rules, etc.

Open questions

- Deeper study of the full Starobinsky model embedded in this non-local setup.
- Explicit computation of the one-loop divergences in this model
- This theory does not a priori prohibits a coexistence of a bounce and inflation. The question is to find such a configuration
- Derive the graviton action from the SFT in the full rigor.

Thank you for listening!