

High-Energy Lorentz Violation, Neutrino Masses And Scalarless Standard Model

Damiano Anselmi

Cargèse – 27/04/2009

based on the papers

[0a] D.A., *Standard Model Without Elementary Scalars And High Energy Lorentz Violation*, [arXiv:0904.1849](#) [[hep-ph](#)]

[0b] D.A., *Weighted power counting, neutrino masses and Lorentz violating extensions of the Standard Model*, [Phys. Rev. D 79 \(2009\) 025017](#) and [arXiv:0808.3475](#) [[hep-ph](#)]

based on the papers

[0a] D.A., *Standard Model Without Elementary Scalars And High Energy Lorentz Violation*, [arXiv:0904.1849](#) [[hep-ph](#)]

[0b] D.A., *Weighted power counting, neutrino masses and Lorentz violating extensions of the Standard Model*, [Phys. Rev. D 79 \(2009\) 025017](#) and [arXiv:0808.3475](#) [[hep-ph](#)]

and previous papers

[1] D.A. and M. Halat, *Renormalization of Lorentz violating theories*, [Phys. Rev. D 76 \(2007\) 125011](#) and [arxiv:0707.2480](#) [[hep-th](#)]

[2] D.A., *Weighted scale invariant quantum field theories*, [JHEP 02 \(2008\) 05](#) and [arxiv:0801.1216](#) [[hep-th](#)]

[3] D.A., *Weighted power counting and Lorentz violating gauge theories. I: General properties*, [Ann. Phys. 324 \(2009\) 874](#) and [arXiv:0808.3470](#) [[hep-th](#)]

[4] D.A., *Weighted power counting and Lorentz violating gauge theories. II: Classification*, [Ann. Phys. 324 \(2009\) 1058](#) and [arXiv:0808.3474](#) [[hep-th](#)]

Lorentz symmetry is a basic ingredient of the Standard Model of particles physics.

Lorentz symmetry is a basic ingredient of the Standard Model of particles physics.

However, several authors have argued that at high energies Lorentz symmetry and possibly CPT could be broken

- [1] V.A. Kostelecký and S. Samuel, Spontaneous breaking of Lorentz symmetry in string theory, Phys. Rev. D 39 (1989) 683;
- S.M. Carroll, G.B. Field and R. Jackiw, Limits on a Lorentz and Parity Violating Modification of Electrodynamics, Phys. Rev. D 41 (1990) 1231;
- G. Amelino-Camelia, J.R. Ellis, N.E. Mavromatos, D.V. Nanopoulos, S. Sarkar, Tests of quantum gravity from observations of gamma-ray bursts, Nature 393 (1998) 763 and arXiv:astro-ph/9712103;
- S.R. Coleman and G.L. Glashow, High-energy tests of Lorentz invariance, Phys. Rev. D 59 (1999) 116008 and arXiv:hep-ph/9812418.
- [2] D. Colladay and V.A. Kostelecký, Lorentz-violating extension of the Standard Model, Phys. Rev. D 58 (1998) 116002 and arXiv:hep-ph/9809521.
- [3] R. Jackiw and V.A. Kostelecký, Radiatively induced Lorentz and CPT violation in electrodynamics, Phys. Rev. Lett. 82 (1999) 3572 and arXiv:hep-ph/9901358;
- O. Bertolami and C.S. Carvalho, Proposed astrophysical test of Lorentz invariance, Phys. Rev. D 61 (2000) 103002 and arXiv:gr-qc/9912117;
- M. Perez-Victoria, Physical (ir)relevance of ambiguities to Lorentz and CPT violation in QED, JHEP 0104 (2001) 032 and arXiv:hep-th/0102021;
- N.R. Bruno, G. Amelino-Camelia and J. Kowalski-Glikman, Deformed boost transformations that saturate at the Planck scale, Phys. Lett. B 522 (2001) 133 and arXiv:hep-th/0107039;

A.A. Andrianov, P. Giacconi and R. Soldati, Lorentz and CPT violations from Chern-Simons modifications of QED, JHEP 0202 (2002) 030 and arXiv:hep-th/0110279;

V.A. Kostelecký, Gravity, Lorentz violation, and the standard model, Phys. Rev. D 69 (2004) 105009 and arXiv:hep-th/0312310;

N. Arkani-Hamed, H.-C. Cheng, M. Luty and J. Thaler, Universal dynamics of spontaneous Lorentz violation and a new spin-dependent inverse-square law force, JHEP 0507 (2005) 029 and arXiv:hep-ph/0407034;

J. Gamboa, J. Lopez-Sarrion and A.P. Polychronakos, Ultraviolet modified photons and anisotropies in the cosmic microwave background radiation, Phys. Lett. B 634 (2006) 471 and arXiv:hep-ph/0510113;

T. Jacobson, S. Liberati and D. Mattingly, Lorentz violation at high energies: concepts, phenomena and astrophysical constraints, Annals Phys. 321 (2006) 150 and arXiv:astro-ph/0505267.

[4] V.A. Kostelecký, Ed., *Proceedings of the Fourth Meeting on CPT and Lorentz Symmetry*, World Scientific, Singapore, 2008;

V.A. Kostelecký and N. Russell, *Data tables for Lorentz and CTP violation*, *ibid.* p. 308 and arXiv:0801.0287 [hep-ph].

- [11] H.B. Nielsen and M. Ninomiya, Beta function in a noncovariant Yang-Mills theory, Nucl.Phys. B 141 (1978) 153;
S. Chadha and H.B. Nielsen, Lorentz invariance as a low-energy phenomenon, Nucl. Phys. B 217 (1983) 125.
- [12] A.G. Cohen and S.L. Glashow, Very special relativity, Phys. Rev. Lett. 97 (2006) 021601 and arXiv:hep-ph/0601236.
- [13] A.G. Cohen and S.L. Glashow, A Lorentz-violating origin of the neutrino mass?, arXiv:hep-ph/0605036.
- [14] V.A. Kostelecký and M. Mewes, Lorentz and CPT violation in the neutrino sector, Phys. Rev. D 70 (2004) 031902 and arXiv:hep-ph/0308300;
V.A. Kostelecký and M. Mewes, Lorentz and CPT violations in neutrinos, Phys. Rev. D 69 (2004) 016005 and arXiv:hep-ph/0309025.
- [15] T. Katori, V.A. Kostelecký and R. Tayloe, Global three-parameter model for neutrino oscillations using Lorentz violation, Phys. Rev. D 74 (2006) 105009 and arXiv:hep-ph/0606154.
- [16] F. Klinkhamer, Lorentz-noninvariant neutrino oscillations: models and predictions, Int. J. Mod. Phys. A 21 (2006) 161 and arXiv:hep-ph/0407200.

Lorentz symmetry is a basic ingredient of the Standard Model of particles physics.

However, several authors have argued that at high energies Lorentz symmetry and possibly CPT could be broken.

The **Lorentz violating parameters** of the Standard Model (Colladay-Kostelecky) extended in the power-counting renormalizable sector have been measured with **great precision**.

Lorentz symmetry is a basic ingredient of the Standard Model of particles physics.

However, several authors have argued that at high energies Lorentz symmetry and possibly CPT could be broken.

The **Lorentz violating parameters** of the Standard Model (Colladay-Kostelecky) extended in the power-counting renormalizable sector have been measured with **great precision**.

It turns out that **Lorentz symmetry** is a very precise symmetry of Nature, at least in low-energy domain.

Lorentz symmetry is a basic ingredient of the Standard Model of particles physics.

However, several authors have argued that at high energies Lorentz symmetry and possibly CPT could be broken.

The **Lorentz violating parameters** of the Standard Model (Colladay-Kostelecky) extended in the power-counting renormalizable sector have been measured with **great precision**.

It turns out that **Lorentz symmetry** is a very precise symmetry of Nature, at least in low-energy domain.

Several (dimensionless) parameters have bounds

$$10^{-15}, \quad 10^{-30}, \quad 10^{-40}$$

V.A. Kostelecký, Ed., *Proceedings of the Fourth Meeting on CPT and Lorentz Symmetry*,
World Scientific, Singapore, 2008;
V.A. Kostelecký and N. Russell, *Data tables for Lorentz and CTP violation*, *ibid.* p. 308 and
arXiv:0801.0287 [hep-ph].

TABLE IV: Electron sector

Combination	Result	System	Ref.
\tilde{b}_X	$(-0.9 \pm 1.4) \times 10^{-31}$ GeV	"	[17]
\tilde{b}_Y	$(-0.9 \pm 1.4) \times 10^{-31}$ GeV	"	[17]
\tilde{b}_Z	$(-0.3 \pm 4.4) \times 10^{-30}$ GeV	"	[17]
$\frac{1}{2}(\tilde{b}_T + \tilde{d}_- - 2\tilde{g}_c - 3\tilde{g}_T + 4\tilde{d}_+ - \tilde{d}_Q)$	$(0.9 \pm 2.2) \times 10^{-27}$ GeV	"	[17]
$\frac{1}{2}(2\tilde{g}_c - \tilde{g}_T - \tilde{b}_T + 4\tilde{d}_+ - \tilde{d}_- - \tilde{d}_Q)$ + $\tan \eta(\tilde{d}_{YZ} - \tilde{H}_{XT})$	$(-0.8 \pm 2.0) \times 10^{-27}$ GeV	"	[17]
\tilde{b}_X	$(0.1 \pm 2.4) \times 10^{-31}$ GeV	"	[18]
\tilde{b}_Y	$(-1.7 \pm 2.5) \times 10^{-31}$ GeV	"	[18]
\tilde{b}_Z	$(-29 \pm 39) \times 10^{-31}$ GeV	"	[18]
\tilde{b}_\perp	$< 3.1 \times 10^{-29}$ GeV	"	[19]
$ \tilde{b}_Z $	$< 7.1 \times 10^{-28}$ GeV	"	[19]
\tilde{b}_X	$(2.8 \pm 6.1) \times 10^{-29}$ GeV	K/He magnetometer	[20]
\tilde{b}_Y	$(6.8 \pm 6.1) \times 10^{-29}$ GeV	"	[20]
r_e	$< 3.2 \times 10^{-24}$	Hg/Cs comparison	[21]
$ \vec{b} $	< 20 radians/s	Penning trap	[22]
$r_{\omega_e^-}$, diurnal	$< 1.6 \times 10^{-21}$	"	[23]
$ \tilde{b}_J $ ($J = X, Y$)	$< 10^{-27}$ GeV	Hg/Cs comparison	[24]*
$c_{XX} - c_{YY}$	$(-2.9 \pm 6.3) \times 10^{-16}$	Optical, microwave resonators	[25]*
$\frac{1}{2}c_{(XY)}$	$(2.1 \pm 0.9) \times 10^{-16}$	"	[25]*
$\frac{1}{2}c_{(XZ)}$	$(-1.5 \pm 0.9) \times 10^{-16}$	"	[25]*
$\frac{1}{2}c_{(YZ)}$	$(-0.5 \pm 1.2) \times 10^{-16}$	"	[25]*
$c_{XX} + c_{YY} - 2c_{ZZ}$	$(-106 \pm 147) \times 10^{-16}$	"	[25]*
λ^{ZZ}	$(13.3 \pm 9.8) \times 10^{-16}$	"	[25]*
$c_{(YZ)}$	$(2.1 \pm 4.6) \times 10^{-16}$	"	[26]*
$c_{(XZ)}$	$(-1.6 \pm 6.3) \times 10^{-16}$	"	[26]*
$c_{(XY)}$	$(7.6 \pm 3.5) \times 10^{-16}$	"	[26]*
$c_{XX} - c_{YY}$	$(1.15 \pm 0.64) \times 10^{-15}$	"	[26]*
$ c_{XX} + c_{YY} - 2c_{ZZ} - 0.25(\tilde{\kappa}_{e-})^{ZZ} $	$< 10^{-12}$	"	[26]*
$ \frac{1}{2}c_{(XY)} $	$< 8 \times 10^{-15}$	Optical resonators	[27]*
$ c_{XX} - c_{YY} $	$< 1.6 \times 10^{-15}$	"	[27]*
$ c_{XX} + c_{YY} - 2c_{ZZ} $	$< 10^{-5}$	Heavy-ion storage ring	[28]*
$ c_{(TX)} , c_{(TY)} , c_{(TZ)} $	$< 10^{-2}$	"	[28]*

V.A. Kostelecký, Ed., *Proceedings of the Fourth Meeting on CPT and Lorentz Symmetry*,
World Scientific, Singapore, 2008;
V.A. Kostelecký and N. Russell, *Data tables for Lorentz and CTP violation*, *ibid.* p. 308 and
arXiv:0801.0287 [hep-ph].

TABLE VII: Photon sector

Combination	Result	System	Ref.
$(\tilde{\kappa}_{e-})^{XY}$	$(-0.1 \pm 0.6) \times 10^{-17}$	Rotating optical resonators	[42]
"	$(-7.7 \pm 4.0) \times 10^{-16}$	Optical, microwave resonators	[25]*
"	$(2.9 \pm 2.3) \times 10^{-16}$	Rotating microwave resonators	[43]
"	$(-3.1 \pm 2.5) \times 10^{-16}$	Rotating optical resonators	[44]
"	$(-0.63 \pm 0.43) \times 10^{-15}$	Rotating microwave resonators	[45]
"	$(-1.7 \pm 1.6) \times 10^{-15}$	Optical, microwave resonators	[26]*
"	$(-5.7 \pm 2.3) \times 10^{-15}$	Microwave resonator, maser	[46]
"	$(1.7 \pm 2.6) \times 10^{-15}$	Optical resonators	[47]
"	$(1.4 \pm 1.4) \times 10^{-13}$	Microwave resonators	[48]
$(\tilde{\kappa}_{e-})^{XZ}$	$(-2.0 \pm 0.9) \times 10^{-17}$	Rotating optical resonators	[42]
"	$(-10.3 \pm 3.9) \times 10^{-16}$	Optical, microwave resonators	[25]*
"	$(-6.9 \pm 2.2) \times 10^{-16}$	Rotating microwave resonators	[43]
"	$(5.7 \pm 4.9) \times 10^{-16}$	Rotating optical resonators	[44]
"	$(0.19 \pm 0.37) \times 10^{-15}$	Rotating microwave resonators	[45]
"	$(-4.0 \pm 3.3) \times 10^{-15}$	Optical, microwave resonators	[26]*
"	$(-3.2 \pm 1.3) \times 10^{-15}$	Microwave resonator, maser	[46]
"	$(-6.3 \pm 12.4) \times 10^{-15}$	Optical resonators	[47]
"	$(-3.5 \pm 4.3) \times 10^{-13}$	Microwave resonators	[48]
$(\tilde{\kappa}_{e-})^{YZ}$	$(-0.3 \pm 1.4) \times 10^{-17}$	Rotating optical resonators	[42]
"	$(0.9 \pm 4.2) \times 10^{-16}$	Optical, microwave resonators	[25]*
"	$(2.1 \pm 2.1) \times 10^{-16}$	Rotating microwave resonators	[43]
"	$(-1.5 \pm 4.4) \times 10^{-16}$	Rotating optical resonators	[44]
"	$(-0.45 \pm 0.37) \times 10^{-15}$	Rotating microwave resonators	[45]
"	$(0.52 \pm 2.52) \times 10^{-15}$	Optical, microwave resonators	[26]*
"	$(-0.5 \pm 1.3) \times 10^{-15}$	Microwave resonator, maser	[46]
"	$(3.6 \pm 9.0) \times 10^{-15}$	Optical resonators	[47]
"	$(1.7 \pm 3.6) \times 10^{-13}$	Microwave resonators	[48]

TABLE VII: Photon sector (continued)

Combination	Result	System	Ref.
$\tilde{\kappa}_{\text{tr}} - \frac{4}{3}C_{00}^e$	$(-5.8 \text{ to } 12) \times 10^{-12}$	Collider physics	[51]*
$\tilde{\kappa}_{\text{tr}} - \frac{4}{3}C_{00}^\nu$	$< 6 \times 10^{-20}$	Astrophysics	[52]*
$-\left[\tilde{\kappa}_{\text{tr}} - \frac{4}{3}C_{00}^e\right]$	$< 9 \times 10^{-16}$	"	[52]*
$\tilde{\kappa}_{\text{tr}}$	$< 1.4 \times 10^{-19}$	"	[50]*
$ \tilde{\kappa}_{\text{tr}} $	$< 8.4 \times 10^{-8}$	Optical atomic clocks	[53]
"	$< 2.2 \times 10^{-7}$	Heavy-ion storage ring	[54]*
"	$< 2 \times 10^{-14}$	Astrophysics	[55]*
"	$< 3 \times 10^{-8}$	$g_e - 2$	[55]*
"	$< 1.6 \times 10^{-5}$	Sagnac interferometer	[56]*
$k_{(E)20}^{(4)}$	$\pm(17_{-9}^{+7}) \times 10^{-31}$	CMB polarization	[10]*
$k_{(B)20}^{(4)}$	$+(17_{-9}^{+7}) \times 10^{-31}$	"	[10]*
$ k^a $ for some a	$< 2 \times 10^{-37}$	Cosmological birefringence	[57]*
k^a for $a = 1, \dots, 10$	$< 2 \times 10^{-32}$	"	[9]*
$k_{(V)10}^{(3)}$	$< 16 \times 10^{-21}$ GeV	Schumann resonances	[58]*
$k_{(V)11}^{(3)}$	$< 12 \times 10^{-21}$ GeV	"	[58]*
$\left(6 k_{(V)11}^{(3)} ^2 + 3 k_{(V)10}^{(3)} ^2\right)^{1/2} / \sqrt{4\pi}$	$(10_{-8}^{+4}) \times 10^{-43}$ GeV	CMB polarization	[12]*
$k_{(V)10}^{(3)}$	$\pm(3 \pm 1) \times 10^{-42}$ GeV	"	[10]*
$k_{(V)11}^{(3)}$	$-(21_{-9}^{+7}) \times 10^{-43}$ GeV	"	[10]*
$k_{(V)00}^{(3)}$	$< 14 \times 10^{-21}$ GeV	Schumann resonances	[58]*
$k_{(V)00}^{(3)}$	$(-1.4 + 0.9 + 0.5) \times 10^{-43}$ GeV	CMB polarization	[59]
"	$(2.3 \pm 5.4) \times 10^{-43}$ GeV	"	[12]*
"	$< 2.5 \times 10^{-43}$ GeV	"	[60]*, [12]*
"	$(1.2 \pm 2.2) \times 10^{-43}$ GeV	"	[61], [12]*
"	$(12 \pm 7) \times 10^{-43}$ GeV	"	[10]*
"	$(2.6 \pm 1.9) \times 10^{-43}$ GeV	"	[62]*, [12]*
"	$(2.5 \pm 3.0) \times 10^{-43}$ GeV	"	[63]*, [12]*
"	$(6.0 \pm 4.0) \times 10^{-43}$ GeV	"	[64]*, [10]*
"	$< 4 \times 10^{-42}$ GeV	Cosmological birefringence	[65]*, [10]*

V.A. Kostelecký, Ed., *Proceedings of the Fourth Meeting on CPT and Lorentz Symmetry*,
World Scientific, Singapore, 2008;
V.A. Kostelecký and N. Russell, *Data tables for Lorentz and CTP violation*, *ibid.* p. 308 and
arXiv:0801.0287 [hep-ph].

Recent analysis of gamma-ray bursts suggests a bound on the scale \bar{M} of Lorentz violation in the dispersion relation

$$c(E) \sim c \left(1 - \frac{E}{\bar{M}} \right)$$

Recent analysis of gamma-ray bursts suggests a bound on the scale \bar{M} of Lorentz violation in the dispersion relation

$$c(E) \sim c \left(1 - \frac{E}{\bar{M}} \right)$$

and found

$$\bar{M} \geq 1.3 \cdot 10^{18} \text{ GeV}$$

Recent analysis of gamma-ray bursts suggests a bound on the scale \bar{M} of Lorentz violation in the dispersion relation

$$c(E) \sim c \left(1 - \frac{E}{\bar{M}} \right)$$

and found

$$\bar{M} \geq 1.3 \cdot 10^{18} \text{ GeV}$$

Since in local QFT odd powers of the energy mean that CPT is violated also, we can take this bound to be a bound on the scale Λ_{CPT} of CPT violation.

Recent analysis of gamma-ray bursts suggests a bound on the scale \bar{M} of Lorentz violation in the dispersion relation

$$c(E) \sim c \left(1 - \frac{E}{\bar{M}} \right)$$

and found

$$\bar{M} \geq 1.3 \cdot 10^{18} \text{ GeV}$$

Since in local QFT odd powers of the energy mean that CPT is violated also, we can take this bound to be a bound on the scale Λ_{CPT} of CPT violation.

On the other hand, Lorentz violation does not necessarily imply CPT violation, so We may assume that there exist two scales, one scale Λ_L for the Lorentz Violation, and one scale Λ_{CPT} for the CPT violation, with

$$\Lambda_{\text{CPT}} \geq \Lambda_L$$

For the same reason, we may assume that there exists an energy range

$$\Lambda_L \leq E \leq \Lambda_{\text{CPT}}$$

that is well described by a Lorentz violating, but CPT invariant quantum field theory.

For the same reason, we may assume that there exists an energy range

$$\Lambda_L \leq E \leq \Lambda_{\text{CPT}}$$

that is well described by a Lorentz violating, but CPT invariant quantum field theory.

Our estimate will give

$$\Lambda_L \sim 10^{14} \text{GeV}$$

so the mentioned range spans **at least 4-5 orders of magnitude**.

For the same reason, we may assume that there exists an energy range

$$\Lambda_L \leq E \leq \Lambda_{\text{CPT}}$$

that is well described by a Lorentz violating, but CPT invariant quantum field theory.

Our estimate will give

$$\Lambda_L \sim 10^{14} \text{GeV}$$

so the mentioned range spans **at least 4-5 orders of magnitude**.

Why is it interesting to consider quantum field theories where Lorentz symmetry
Is **explicitly** broken?

The set of power-counting renormalizable theories is considerably “small”

The set of power-counting renormalizable theories is considerably “small”

Relaxing some assumptions can enlarge it, but often it enlarges it too much

The set of power-counting renormalizable theories is considerably “small”

Relaxing some assumptions can enlarge it, but often it enlarges it too much

Without **locality** in principle every theory can be made finite

The set of power-counting renormalizable theories is considerably “small”

Relaxing some assumptions can enlarge it, but often it enlarges it too much

Without **locality** in principle every theory can be made finite

Without **unitarity** even gravity can be renormalized

The set of power-counting renormalizable theories is considerably “small”

Relaxing some assumptions can enlarge it, but often it enlarges it too much

Without **locality** in principle every theory can be made finite

Without **unitarity** even gravity can be renormalized

Relaxing **Lorentz invariance** appears to be interesting in its own right

The set of power-counting renormalizable theories is considerably “small”

Relaxing some assumptions can enlarge it, but often it enlarges it too much

Without **locality** in principle every theory can be made finite

Without **unitarity** even gravity can be renormalized

Relaxing **Lorentz invariance** appears to be interesting in its own right

It could be useful to define the **ultraviolet limit of quantum gravity**, to study **extensions of the Standard Model**, **effective field theories**, nuclear physics, and the theory of critical phenomena

The set of power-counting renormalizable theories is considerably “small”

Relaxing some assumptions can enlarge it, but often it enlarges it too much

Without **locality** in principle every theory can be made finite

Without **unitarity** even gravity can be renormalized

Relaxing **Lorentz invariance** appears to be interesting in its own right

It could be useful to define the **ultraviolet limit of quantum gravity**, to study **extensions of the Standard Model**, **effective field theories**, nuclear physics, and the theory of critical phenomena

Here we are interested in the renormalization of Lorentz violating theories obtained improving the behavior of propagators with the help of **higher space derivatives**

The set of power-counting renormalizable theories is considerably “small”

Relaxing some assumptions can enlarge it, but often it enlarges it too much

Without **locality** in principle every theory can be made finite

Without **unitarity** even gravity can be renormalized

Relaxing **Lorentz invariance** appears to be interesting in its own right

It could be useful to define the **ultraviolet limit of quantum gravity**, to study **extensions of the Standard Model**, **effective field theories**, nuclear physics, and the theory of critical phenomena

Here we are interested in the renormalization of Lorentz violating theories obtained improving the behavior of propagators with the help of

higher space derivatives

and study under which conditions

no higher time derivatives are turned on

The set of power-counting renormalizable theories is considerably “small”

Relaxing some assumptions can enlarge it, but often it enlarges it too much

Without **locality** in principle every theory can be made finite

Without **unitarity** even gravity can be renormalized

Relaxing **Lorentz invariance** appears to be interesting in its own right

It could be useful to define the **ultraviolet limit of quantum gravity**, to study **extensions of the Standard Model**, **effective field theories**, nuclear physics, and the theory of critical phenomena

Here we are interested in the renormalization of Lorentz violating theories obtained improving the behavior of propagators with the help of

higher space derivatives

and study under which conditions

no higher time derivatives are turned on

to be consistent with **unitarity**

The set of power-counting renormalizable theories is considerably “small”

Relaxing some assumptions can enlarge it, but often it enlarges it too much

Without **locality** in principle every theory can be made finite

Without **unitarity** even gravity can be renormalized

Relaxing **Lorentz invariance** appears to be interesting in its own right

It could be useful to define the **ultraviolet limit of quantum gravity**, to study **extensions of the Standard Model**, **effective field theories**, nuclear physics, and the theory of critical phenomena

Here we are interested in the renormalization of Lorentz violating theories obtained improving the behavior of propagators with the help of

higher space derivatives

and study under which conditions

no higher time derivatives are turned on

to be consistent with **unitarity**

The approach that I proposed is based of a **modified** criterion of power counting, dubbed **weighted power counting**

Scalar fields

Scalar fields

Break spacetime in two pieces:

$$M_D = M_{\hat{D}} \otimes M_{\overline{D}}$$

$$d = \hat{d} + \overline{d}$$

Scalar fields

Break spacetime in two pieces:

$$M_D = M_{\hat{D}} \otimes M_{\bar{D}} \qquad d = \hat{d} + \bar{d}$$

Break coordinates and momenta correspondingly: $p = (\hat{p}, \bar{p})$

Scalar fields

Break spacetime in two pieces:

$$M_D = M_{\hat{D}} \otimes M_{\bar{D}} \qquad d = \hat{d} + \bar{d}$$

Break coordinates and momenta correspondingly: $p = (\hat{p}, \bar{p})$

Consider
the free theory

$$\mathcal{L}_{\text{free}} = \frac{1}{2}(\hat{\partial}\varphi)^2 + \frac{1}{2\Lambda_L^{2n-2}}(\bar{\partial}^n \varphi)^2.$$

Scalar fields

Break spacetime in two pieces:

$$M_D = M_{\hat{D}} \otimes M_{\bar{D}} \qquad d = \hat{d} + \bar{d}$$

Break coordinates and momenta correspondingly: $p = (\hat{p}, \bar{p})$

Consider
the free theory

$$\mathcal{L}_{\text{free}} = \frac{1}{2}(\hat{\partial}\varphi)^2 + \frac{1}{2\Lambda_L^{2n-2}}(\bar{\partial}^n \varphi)^2.$$

This free theory is invariant under the “weighted” scale transformation

$$\hat{x} \rightarrow \hat{x} e^{-\Omega}, \qquad \bar{x} \rightarrow \bar{x} e^{-\Omega/n}, \qquad \varphi \rightarrow \varphi e^{\Omega(\hat{d}/2-1)}$$

where $\hat{d} \equiv \hat{d} + \bar{d}/n$ is the “weighted dimension”

The propagator

$$\frac{1}{\hat{k}^2 + \frac{(\bar{k}^2)^n}{\Lambda_L^{2(n-1)}}}$$

behaves better than usual in the barred directions

The propagator

$$\frac{1}{\widehat{k}^2 + \frac{(\overline{k}^2)^n}{\Lambda_L^{2(n-1)}}}$$

behaves better than usual in the barred directions

Adding “weighted relevant” terms we get a free theory

$$\frac{1}{2}(\widehat{\partial}\varphi)^2 + \sum_{k=0}^n, \frac{\lambda_k M^{2(1-k/n)}}{2\Lambda_L^{2k(n-1)/n}} (\overline{\partial}^k \varphi)^2$$

The propagator

$$\frac{1}{\widehat{k}^2 + \frac{(\overline{k}^2)^n}{\Lambda_L^{2(n-1)}}}$$

behaves better than usual in the barred directions

Adding “weighted relevant” terms we get a free theory

$$\frac{1}{2}(\widehat{\partial}\varphi)^2 + \sum_{k=0}^n \frac{\lambda_k M^{2(1-k/n)}}{2\Lambda_L^{2k(n-1)/n}} (\overline{\partial}^k \varphi)^2$$

that flows to the previous one in the UV and to the Lorentz invariant free theory in the infrared

(actually, the IR Lorentz recovery is much more subtle, see below)

Add **vertices** $\left[\hat{\partial}^{p_1} \bar{\partial}^{p_2} \varphi^N \right]_{\alpha}$ constructed with φ , $\hat{\partial}$ and $\bar{\partial}$.

Call $\delta_N^{(\alpha)} = p_1 + \frac{p_2}{n}$ their degrees under

$$\hat{x} \rightarrow \hat{x} e^{-\Omega}, \quad \bar{x} \rightarrow \bar{x} e^{-\Omega/n}$$

N = number of legs, α = extra label

Add **vertices** $\left[\hat{\partial}^{p_1} \bar{\partial}^{p_2} \varphi^N \right]_{\alpha}$ constructed with φ , $\hat{\partial}$ and $\bar{\partial}$.

Call $\delta_N^{(\alpha)} = p_1 + \frac{p_2}{n}$ their degrees under

$$\hat{x} \rightarrow \hat{x} e^{-\Omega}, \quad \bar{x} \rightarrow \bar{x} e^{-\Omega/n}$$

N = number of legs, α = extra label

Other quadratic terms can be treated as “vertices” for the purposes of renormalization

$$\frac{a_m}{2\Lambda_L^{2m-2}} (\bar{\partial}^m \varphi)^2, \quad m < n,$$

Consider a diagram G with L loops, I internal legs, E external legs and $v_N^{(\alpha)}$ vertices of type (N, α)

Consider a diagram G with L loops, I internal legs, E external legs and $v_N^{(\alpha)}$ vertices of type (N, α)

$$\mathcal{I}_G(k) = \int \frac{d^{L\hat{D}}\hat{p}}{(2\pi)^{L\hat{D}}} \int \frac{d^{L\bar{D}}\bar{p}}{(2\pi)^{L\bar{D}}} \prod_{i=1}^I \mathcal{P}_{-2,n}^{(i)}(p, k) \prod_{j=1}^V \mathcal{V}_{\delta_j, n}^{(j)}(p, k).$$

Consider a diagram G with L loops, I internal legs, E external legs and $v_N^{(\alpha)}$ vertices of type (N, α)

$$\mathcal{I}_G(k) = \int \frac{d^{L\hat{D}}\hat{p}}{(2\pi)^{L\hat{D}}} \int \frac{d^{L\overline{D}}\overline{p}}{(2\pi)^{L\overline{D}}} \prod_{i=1}^I \mathcal{P}_{-2,n}^{(i)}(p, k) \prod_{j=1}^V \mathcal{V}_{\delta_j, n}^{(j)}(p, k),$$

$d^{\hat{D}}\hat{p} d^{\overline{D}}\overline{p}$ is a **weighted measure** of degree $D \equiv \hat{D} + \overline{D}/n$

Consider a diagram G with L loops, I internal legs, E external legs and $v_N^{(\alpha)}$ vertices of type (N, α)

$$\mathcal{I}_G(k) = \int \frac{d^{L\hat{D}}\hat{p}}{(2\pi)^{L\hat{D}}} \int \frac{d^{L\overline{D}}\overline{p}}{(2\pi)^{L\overline{D}}} \prod_{i=1}^I \mathcal{P}_{-2,n}^{(i)}(p, k) \prod_{j=1}^V \mathcal{V}_{\delta_j, n}^{(j)}(p, k).$$

$d^{\hat{D}}\hat{p} d^{\overline{D}}\overline{p}$ is a **weighted measure** of degree $D \equiv \hat{D} + \overline{D}/n$

Performing a “**weighted rescaling**” $(\hat{k}, \bar{k}) \rightarrow (\lambda\hat{k}, \lambda^{1/n}\bar{k})$ of external momenta, together with a change of variables $(\hat{p}, \overline{p}) \rightarrow (\lambda\hat{p}, \lambda^{1/n}\overline{p})$

Consider a diagram G with L loops, I internal legs, E external legs and $v_N^{(\alpha)}$ vertices of type (N, α)

$$\mathcal{I}_G(k) = \int \frac{d^{L\hat{D}}\hat{p}}{(2\pi)^{L\hat{D}}} \int \frac{d^{L\overline{D}}\overline{p}}{(2\pi)^{L\overline{D}}} \prod_{i=1}^I \mathcal{P}_{-2,n}^{(i)}(p, k) \prod_{j=1}^V \mathcal{V}_{\delta_j, n}^{(j)}(p, k).$$

$d^{\hat{D}}\hat{p} d^{\overline{D}}\overline{p}$ is a **weighted measure** of degree $\mathcal{D} \equiv \hat{D} + \overline{D}/n$

Performing a “**weighted rescaling**” $(\hat{k}, \bar{k}) \rightarrow (\lambda\hat{k}, \lambda^{1/n}\bar{k})$ of external momenta, together with a change of variables $(\hat{p}, \overline{p}) \rightarrow (\lambda\hat{p}, \lambda^{1/n}\overline{p})$

we see that $\mathcal{I}_G(k)$ is a **homogeneous weighted function** of degree

$$L\mathcal{D} - 2I + \sum_{j=1}^V \delta_j = L\mathcal{D} - 2I + \sum_{(N, \alpha)} \delta_N^{(\alpha)} v_N^{(\alpha)}$$

Consider a diagram G with L loops, I internal legs, E external legs and $v_N^{(\alpha)}$ vertices of type (N, α)

$$\mathcal{I}_G(k) = \int \frac{d^{L\hat{D}}\hat{p}}{(2\pi)^{L\hat{D}}} \int \frac{d^{L\overline{D}}\overline{p}}{(2\pi)^{L\overline{D}}} \prod_{i=1}^I \mathcal{P}_{-2,n}^{(i)}(p, k) \prod_{j=1}^V \mathcal{V}_{\delta_j, n}^{(j)}(p, k).$$

$d^{\hat{D}}\hat{p} d^{\overline{D}}\overline{p}$ is a **weighted measure** of degree $\mathcal{D} \equiv \hat{D} + \overline{D}/n$

Performing a “**weighted rescaling**” $(\hat{k}, \bar{k}) \rightarrow (\lambda\hat{k}, \lambda^{1/n}\bar{k})$ of external momenta, together with a change of variables $(\hat{p}, \overline{p}) \rightarrow (\lambda\hat{p}, \lambda^{1/n}\overline{p})$

we see that $\mathcal{I}_G(k)$ is a **homogeneous weighted function** of degree

$$L\mathcal{D} - 2I + \sum_{j=1}^V \delta_j = L\mathcal{D} - 2I + \sum_{(N,\alpha)} \delta_N^{(\alpha)} v_N^{(\alpha)}$$

Its overall divergent part is a **homogeneous weighted polynomial** of degree

$$\omega(G) = L\mathcal{D} - 2I + \sum_{(N,\alpha)} \delta_N^{(\alpha)} v_N^{(\alpha)}$$

Using the standard relations

$$L = I - V + 1, \quad E + 2I = \sum_{(N, \alpha)} N v_N^{(\alpha)}$$

Using the standard relations

$$L = I - V + 1, \quad E + 2I = \sum_{(N,\alpha)} N v_N^{(\alpha)}$$

we get

$$\omega(G) = d(E) + \sum_{(N,\alpha)} v_N^{(\alpha)} \left[\delta_N^{(\alpha)} - d(N) \right]$$

Where

$$d(X) \equiv d \left(1 - \frac{X}{2} \right) + X$$

Using the standard relations

$$L = I - V + 1, \quad E + 2I = \sum_{(N,\alpha)} N v_N^{(\alpha)}$$

we get

$$\omega(G) = d(E) + \sum_{(N,\alpha)} v_N^{(\alpha)} \left[\delta_N^{(\alpha)} - d(N) \right]$$

Where

$$d(X) \equiv d \left(1 - \frac{X}{2} \right) + X$$

Renormalizable theories have

$$\delta_N^{(\alpha)} \leq d(N)$$

Using the standard relations

$$L = I - V + 1, \quad E + 2I = \sum_{(N,\alpha)} N v_N^{(\alpha)}$$

we get

$$\omega(G) = d(E) + \sum_{(N,\alpha)} v_N^{(\alpha)} \left[\delta_N^{(\alpha)} - d(N) \right]$$

Where

$$d(X) \equiv d \left(1 - \frac{X}{2} \right) + X$$

Renormalizable theories have

$$\delta_N^{(\alpha)} \leq d(N)$$

Indeed $\delta_N^{(\alpha)} \leq d(N)$ implies $\omega(G) \leq d(E)$

Writing

$$d(N) = d - \frac{N}{2} (d - 2)$$

Writing

$$d(N) = \vec{d} - \frac{N}{2} (\vec{d} - 2)$$

we see that [polynomiality](#) demands $\vec{d} > 2$

Writing

$$d(N) = d - \frac{N}{2} (d - 2)$$

we see that **polynomiality** demands $d > 2$

and the **maximal number of legs** is $N_{\max} = \left\lceil \frac{2d}{d-2} \right\rceil$

$E = 2$ implies $\omega(G) \leq 2$

$E > 2$ implies $\omega(G) < 2$

Writing

$$d(N) = \vec{d} - \frac{N}{2} (\vec{d} - 2)$$

we see that **polynomiality** demands $\vec{d} > 2$

and the **maximal number of legs** is $N_{\max} = \left\lceil \frac{2\vec{d}}{\vec{d} - 2} \right\rceil$

$E = 2$ implies $\omega(G) \leq 2$

$E > 2$ implies $\omega(G) < 2$

Conclusion:

**renormalization does not turn on
higher time derivatives**

Examples

$$\vec{d} = 4$$

$$\mathcal{L}_{(2,2n)} = \frac{1}{2}(\widehat{\partial}\varphi)^2 + \frac{1}{2\Lambda_L^{2(n-1)}}(\overline{\partial}^n\varphi)^2 + \frac{\lambda}{4!\Lambda_L^{2(n-1)}}\varphi^4$$

Examples

$$\vec{d} = 4$$

$$\mathcal{L}_{(2,2n)} = \frac{1}{2}(\widehat{\partial}\varphi)^2 + \frac{1}{2\Lambda_L^{2(n-1)}}(\overline{\partial}^n\varphi)^2 + \frac{\lambda}{4!\Lambda_L^{2(n-1)}}\varphi^4$$

n=2: six-dimensional φ^4 -theory

Examples

$$\vec{d} = 4$$

$$\mathcal{L}_{(2,2n)} = \frac{1}{2}(\hat{\partial}\varphi)^2 + \frac{1}{2\Lambda_L^{2(n-1)}}(\bar{\partial}^n\varphi)^2 + \frac{\lambda}{4!\Lambda_L^{2(n-1)}}\varphi^4$$

n=2: six-dimensional φ^4 -theory

Strictly-renormalizable models are classically weighted scale invariant, namely invariant under

$$\hat{x} \rightarrow \hat{x} e^{-\Omega}, \quad \bar{x} \rightarrow \bar{x} e^{-\Omega/n}, \quad \varphi \rightarrow \varphi e^{\Omega(\vec{d}/2-1)}$$

Examples

$$\vec{d} = 4$$

$$\mathcal{L}_{(2,2n)} = \frac{1}{2}(\hat{\partial}\varphi)^2 + \frac{1}{2\Lambda_L^{2(n-1)}}(\bar{\partial}^n\varphi)^2 + \frac{\lambda}{4!\Lambda_L^{2(n-1)}}\varphi^4$$

n=2: six-dimensional φ^4 -theory

Strictly-renormalizable models are classically weighted scale invariant, namely invariant under

$$\hat{x} \rightarrow \hat{x} e^{-\Omega}, \quad \bar{x} \rightarrow \bar{x} e^{-\Omega/n}, \quad \varphi \rightarrow \varphi e^{\Omega(\vec{d}/2-1)}$$

The weighted scale invariance is **anomalous** at the quantum level

Four dimensional examples

$$\widehat{d} = 1 \quad n = 2$$

$$\mathcal{L}_{(1,3)} = \frac{1}{2}(\widehat{\partial}\varphi)^2 + \frac{1}{2\Lambda_L^2}(\overline{\Delta}\varphi)^2 + \frac{\lambda_6}{6!\Lambda_L^4}\varphi^4(\overline{\partial}\varphi)^2 + \frac{\lambda_{10}}{10!\Lambda_L^6}\varphi^{10}$$

Four dimensional examples

$$\widehat{d} = 1 \quad n = 2$$

$$\mathcal{L}_{(1,3)} = \frac{1}{2}(\widehat{\partial}\varphi)^2 + \frac{1}{2\Lambda_L^2}(\overline{\Delta}\varphi)^2 + \frac{\lambda_6}{6!\Lambda_L^4}\varphi^4(\overline{\partial}\varphi)^2 + \frac{\lambda_{10}}{10!\Lambda_L^6}\varphi^{10}$$

$$\widehat{d} = 2 \quad n = 2$$

$$\mathcal{L}_{(2,2)} = \frac{1}{2}(\widehat{\partial}\varphi)^2 + \frac{1}{2\Lambda_L^2}(\overline{\Delta}\varphi)^2 + \frac{\lambda_4}{4!\Lambda_L^2}\varphi^2(\overline{\partial}\varphi)^2 + \frac{\lambda_6}{6!\Lambda_L^2}\varphi^6$$

Four dimensional examples

$$\widehat{d} = 1 \quad n = 2$$

$$\mathcal{L}_{(1,3)} = \frac{1}{2}(\widehat{\partial}\varphi)^2 + \frac{1}{2\Lambda_L^2}(\overline{\Delta}\varphi)^2 + \frac{\lambda_6}{6!\Lambda_L^4}\varphi^4(\overline{\partial}\varphi)^2 + \frac{\lambda_{10}}{10!\Lambda_L^6}\varphi^{10}$$

$$\widehat{d} = 2 \quad n = 2$$

$$\mathcal{L}_{(2,2)} = \frac{1}{2}(\widehat{\partial}\varphi)^2 + \frac{1}{2\Lambda_L^2}(\overline{\Delta}\varphi)^2 + \frac{\lambda_4}{4!\Lambda_L^2}\varphi^2(\overline{\partial}\varphi)^2 + \frac{\lambda_6}{6!\Lambda_L^2}\varphi^6$$

$$\widehat{d} = 3 \quad n = 3$$

$$\mathcal{L}_{(3,1)}^{\text{even}} = \frac{1}{2}(\widehat{\partial}\varphi)^2 + \frac{1}{2\Lambda_L^4}(\overline{\partial\Delta}\varphi)^2 + \frac{\lambda_4}{4!\Lambda_L^2}\varphi^2(\overline{\partial}\varphi)^2$$

Källen-Lehman representation and unitarity

$$\Delta(x) \equiv \langle 0 | T \varphi(x) \varphi(0) | 0 \rangle = \int_0^\infty ds \int \frac{d^d k}{(2\pi)^d} \frac{i e^{-ik \cdot x} \rho(s, \vec{k}^2)}{\widehat{k}^2 - s + i\varepsilon}$$

$$\text{Im} \left[\frac{i}{\pi} \langle \tilde{\varphi}(-k) \tilde{\varphi}(k) \rangle \right] = \rho(\widehat{k}^2, \vec{k}^2) \geq 0$$

Cutting rules

$$\Delta^\pm(x) \equiv \int \frac{d^d k}{(2\pi)^d} e^{-ik \cdot x} \theta(\pm k_0) \rho(\widehat{k}^2, \vec{k}^2)$$

$$\Delta(x) = \theta(x_0) \Delta^+(x) + \theta(-x_0) \Delta^-(x)$$

$$\Delta^\mp(-x) = \Delta^\pm(x), \quad \Delta^{\pm*}(x) = \Delta^\mp(x), \quad \Delta^*(x) = \theta(x_0) \Delta^-(x) + \theta(-x_0) \Delta^+(x)$$

Causality

Our theories satisfy Bogoliubov's definition of causality

$$\frac{\delta^2 S}{\delta g(x_i) \delta g(x_j)} S^\dagger + \frac{\delta S}{\delta g(x_i)} \frac{\delta S^\dagger}{\delta g(x_j)} = 0 \quad \text{if } x_{i0} < x_{j0}$$

which is a simple consequence of the largest time equation and the cutting rules

For the two-point function this is just the statement

$$\Delta(x) = \Delta^+(x) \quad \text{if } x_0 > 0$$

immediate consequence of

$$\Delta(x) = \theta(x_0) \Delta^+(x) + \theta(-x_0) \Delta^-(x)$$

Fermions

Fermions

The extension to fermions is straightforward. The free lagrangian is

$$\mathcal{L} = \overline{\psi} \hat{\partial} \psi + \frac{1}{\Lambda_L^{n-1}} \overline{\psi} \partial^n \psi,$$

Fermions

The extension to fermions is straightforward. The free lagrangian is

$$\mathcal{L} = \bar{\psi} \hat{\partial} \psi + \frac{1}{\Lambda_L^{n-1}} \bar{\psi} \bar{\partial}^n \psi,$$

An example is the **four fermion theory** with $\bar{d}=2$.

$$\mathcal{L}_{\bar{d}=2} = \bar{\psi} \left(\hat{\partial} + \frac{\bar{\partial}^n}{\Lambda_L^{n-1}} \right) \psi - \frac{\lambda^2}{2\Lambda_L^{d-2}} (\bar{\psi}\psi)^2$$

Fermions

The extension to fermions is straightforward. The free lagrangian is

$$\mathcal{L} = \bar{\psi} \hat{\partial} \psi + \frac{1}{\Lambda_L^{n-1}} \bar{\psi} \bar{\partial}^n \psi,$$

An example is the **four fermion theory** with $\bar{d}=2$.

$$\mathcal{L}_{\bar{d}=2} = \bar{\psi} \left(\hat{\partial} + \frac{\bar{\partial}^n}{\Lambda_L^{n-1}} \right) \psi - \frac{\lambda^2}{2\Lambda_L^{d-2}} (\bar{\psi}\psi)^2$$

An example of four dimensional **scalar-fermion theory** is

$$\begin{aligned} \mathcal{L}_{(2,2)} = & \bar{\psi} \hat{\partial} \psi + \frac{\eta}{\Lambda_L} \bar{\psi} \bar{\Delta} \psi + \frac{1}{2} (\hat{\partial} \varphi)^2 + \frac{1}{2\Lambda_L^2} (\bar{\Delta} \varphi)^2 \\ & + \frac{\lambda_2}{2\Lambda_L} \varphi^2 \bar{\psi} \psi + \frac{\lambda_4}{4!\Lambda_L^2} \varphi^2 (\bar{\partial} \varphi)^2 + \frac{\lambda_6}{6!\Lambda_L^2} \varphi^6 \end{aligned}$$

Gauge fields

Gauge fields

Gauge fields are more tricky. Decompose the gauge field as $A = (\hat{A}, \bar{A})$

Gauge fields

Gauge fields are more tricky. Decompose the gauge field as $A = (\hat{A}, \bar{A})$ and assign weights

$$[g\hat{A}] = [\hat{D}] = 1, \quad [g\bar{A}] = [\bar{D}] = \frac{1}{n},$$

Gauge fields

Gauge fields are more tricky. Decompose the gauge field as $A = (\hat{A}, \bar{A})$ and assign weights

$$[g\hat{A}] = [\hat{D}] = 1, \quad [g\bar{A}] = [\bar{D}] = \frac{1}{n},$$

so that the **covariant derivative** is decomposed consistently,

$$D = (\hat{D}, \bar{D}) = (\hat{\partial} + g\hat{A}, \bar{\partial} + g\bar{A})$$

Gauge fields

Gauge fields are more tricky. Decompose the gauge field as $A = (\hat{A}, \bar{A})$ and assign weights

$$[g\hat{A}] = [\hat{D}] = 1, \quad [g\bar{A}] = [\bar{D}] = \frac{1}{n},$$

so that the **covariant derivative** is decomposed consistently,

$$D = (\hat{D}, \bar{D}) = (\hat{\partial} + g\hat{A}, \bar{\partial} + g\bar{A})$$

The **field strength** is decomposed in three sets of components,

$$\hat{F}_{\mu\nu} \equiv F_{\hat{\mu}\hat{\nu}}, \quad \tilde{F}_{\mu\nu} \equiv F_{\hat{\mu}\bar{\nu}}, \quad \bar{F}_{\mu\nu} \equiv F_{\bar{\mu}\bar{\nu}}.$$

Gauge fields

Gauge fields are more tricky. Decompose the gauge field as $A = (\hat{A}, \bar{A})$ and assign weights

$$[g\hat{A}] = [\hat{D}] = 1, \quad [g\bar{A}] = [\bar{D}] = \frac{1}{n},$$

so that the **covariant derivative** is decomposed consistently,

$$D = (\hat{D}, \bar{D}) = (\hat{\partial} + g\hat{A}, \bar{\partial} + g\bar{A})$$

The **field strength** is decomposed in three sets of components,

$$\hat{F}_{\mu\nu} \equiv F_{\hat{\mu}\hat{\nu}}, \quad \tilde{F}_{\mu\nu} \equiv F_{\hat{\mu}\bar{\nu}}, \quad \bar{F}_{\mu\nu} \equiv F_{\bar{\mu}\bar{\nu}}.$$

The **quadratic lagrangian** reads

$$\mathcal{L}'_Q = \frac{1}{2}F_{\hat{\mu}\bar{\nu}}^2 - \frac{1}{4}F_{\bar{\mu}\bar{\nu}}\tau'(\bar{\Upsilon})F_{\bar{\mu}\bar{\nu}}$$

where $\bar{\Upsilon} \equiv -\bar{D}^2/\Lambda_L^2$ and τ' is a polynomial of degree $n-1$

The **BRST symmetry** is unmodified

$$\begin{aligned} sA_\mu^a &= D_\mu^{ab}C^b = \partial_\mu C^a + gf^{abc}A_\mu^bC^c, & sC^a &= -\frac{g}{2}f^{abc}C^bC^c, \\ s\bar{C}^a &= B^a, & sB^a &= 0, & s\psi^i &= -gT_{ij}^aC^a\psi^j, \end{aligned}$$

The **BRST symmetry** is unmodified

$$\begin{aligned} sA_\mu^a &= D_\mu^{ab}C^b = \partial_\mu C^a + gf^{abc}A_\mu^bC^c, & sC^a &= -\frac{g}{2}f^{abc}C^bC^c, \\ s\bar{C}^a &= B^a, & sB^a &= 0, & s\psi^i &= -gT_{ij}^aC^a\psi^j, \end{aligned}$$

We choose the **gauge-fixing**

$$\mathcal{L}_{\text{gf}} = s\Psi, \quad \Psi = \bar{C}^a \left(-\frac{\lambda}{2}B^a + \mathcal{G}^a \right), \quad \mathcal{G}^a \equiv \hat{\partial} \cdot \hat{A}^a + \zeta(\bar{v}) \bar{\partial} \cdot \bar{A}^a,$$

where $\bar{v} \equiv -\bar{\partial}^2/\Lambda_L^2$ and ζ is a polynomial of degree $n-1$

The **BRST symmetry** is unmodified

$$\begin{aligned} sA_\mu^a &= D_\mu^{ab}C^b = \partial_\mu C^a + gf^{abc}A_\mu^b C^c, & sC^a &= -\frac{g}{2}f^{abc}C^b C^c, \\ s\bar{C}^a &= B^a, & sB^a &= 0, & s\psi^i &= -gT_{ij}^a C^a \psi^j, \end{aligned}$$

We choose the **gauge-fixing**

$$\mathcal{L}_{\text{gf}} = s\Psi, \quad \Psi = \bar{C}^a \left(-\frac{\lambda}{2} B^a + \mathcal{G}^a \right), \quad \mathcal{G}^a \equiv \hat{\partial} \cdot \hat{A}^a + \zeta(\bar{v}) \bar{\partial} \cdot \bar{A}^a,$$

where $\bar{v} \equiv -\bar{\partial}^2/\Lambda_L^2$ and ζ is a polynomial of degree $n-1$

The **ghost propagator** is

$$\frac{1}{D(1, \zeta)}$$

where

$$D(x, y) \equiv x\hat{k}^2 + y\bar{k}^2$$

The **gauge-field propagator** is involved, but in the “Feynman” gauge

$$\zeta = \lambda = \tau'$$

The **gauge-field propagator** is involved, but in the “Feynman” gauge

$$\zeta = \lambda = \tau'$$

It becomes diagonal and simple:

$$\begin{pmatrix} \langle \hat{A} \hat{A} \rangle & \langle \hat{A} \bar{A} \rangle \\ \langle \bar{A} \hat{A} \rangle & \langle \bar{A} \bar{A} \rangle \end{pmatrix} = \frac{1}{D(1, \tau')} \begin{pmatrix} \tau' & 0 \\ 0 & \bar{\delta} \end{pmatrix}$$

$$D(x, y) \equiv x \hat{k}^2 + y \bar{k}^2$$

The **gauge-field propagator** is involved, but in the “Feynman” gauge

$$\zeta = \lambda = \tau'$$

It becomes diagonal and simple:

$$\begin{pmatrix} \langle \hat{A} \hat{A} \rangle & \langle \hat{A} \bar{A} \rangle \\ \langle \bar{A} \hat{A} \rangle & \langle \bar{A} \bar{A} \rangle \end{pmatrix} = \frac{1}{D(1, \tau')} \begin{pmatrix} \tau' & 0 \\ 0 & \bar{\delta} \end{pmatrix}$$

$$D(x, y) \equiv x \hat{k}^2 + y \bar{k}^2$$

The weight assignments are

$$[\hat{A}] = \frac{\bar{d}}{2} - \frac{1}{n}, \quad [\bar{A}] = \frac{\bar{d}}{2} - 1, \quad [\tilde{F}] = \frac{\bar{d}}{2}, \quad [\bar{F}] = \frac{\bar{d}}{2} - 1 + \frac{1}{n}.$$

$$[g] = 1 + \frac{1}{n} - \frac{\bar{d}}{2}$$

$$[C] = [\bar{C}] = \frac{\bar{d}}{2} - 1, \quad [s] = \frac{1}{n}, \quad [B] = \frac{\bar{d}}{2} - 1 + \frac{1}{n}.$$

The physical degrees of freedom can be read in the Coulomb gauge-fixing

$$\mathcal{G}_C^a = \bar{\partial} \cdot \bar{A}^a.$$

which can be reached taking a suitable limit on the general one.

The **physical degrees of freedom** can be read in the **Coulomb gauge-fixing**

$$\mathcal{G}_C^a = \bar{\partial} \cdot \bar{A}^a.$$

which can be reached taking a suitable limit on the general one.

We get non-propagating ghosts and a quadratic lagrangian

$$\mathcal{L}'_Q + \frac{1}{2\varsigma} (\bar{\partial} \cdot \bar{A}^a)^2$$

which gives the propagator

$$\begin{aligned} \langle \hat{A}(k) \hat{A}(-k) \rangle &= \frac{1}{\bar{k}^2} + \frac{\varsigma \hat{k}^2}{(\bar{k}^2)^2}, & \langle \hat{A}(k) \bar{A}(-k) \rangle &= \frac{\varsigma \hat{k} \bar{k}}{(\bar{k}^2)^2}, \\ \langle \bar{A}(k) \bar{A}(-k) \rangle &= \frac{1}{D(1, \tau')} \left(\bar{\delta} - \frac{\bar{k} \bar{k}}{\bar{k}^2} \right) + \frac{\varsigma \bar{k} \bar{k}}{(\bar{k}^2)^2}. \end{aligned}$$

The **physical degrees of freedom** can be read in the **Coulomb gauge-fixing**

$$\mathcal{G}_C^a = \bar{\partial} \cdot \bar{A}^a.$$

which can be reached taking a suitable limit on the general one.

We get non-propagating ghosts and a quadratic lagrangian

$$\mathcal{L}'_Q + \frac{1}{2\varsigma} (\bar{\partial} \cdot \bar{A}^a)^2$$

which gives the propagator

$$\begin{aligned} \langle \hat{A}(k) \hat{A}(-k) \rangle &= \frac{1}{\bar{k}^2} + \frac{\varsigma \hat{k}^2}{(\bar{k}^2)^2}, & \langle \hat{A}(k) \bar{A}(-k) \rangle &= \frac{\varsigma \hat{k} \bar{k}}{(\bar{k}^2)^2}, \\ \langle \bar{A}(k) \bar{A}(-k) \rangle &= \frac{1}{D(1, \tau')} \left(\bar{\delta} - \frac{\bar{k} \bar{k}}{\bar{k}^2} \right) + \frac{\varsigma \bar{k} \bar{k}}{(\bar{k}^2)^2}. \end{aligned}$$

We have $\bar{d} - 1$ **degrees of freedom** with energies

$$E = \sqrt{\bar{k}^2 \tau' (\bar{k}^2 / \Lambda_L^2)}.$$

A property of gauge theories is that in four dimensions gauge interactions
are always super-renormalizable
from the weighted power-counting viewpoint

A property of gauge theories is that in four dimensions gauge interactions
are always super-renormalizable

from the weighted power-counting viewpoint

Indeed, for

$$\bar{d} = 1 + \frac{3}{n}$$

the weight of the gauge coupling is

$$[g] = \frac{1}{2} - \frac{1}{2n} > 0 \quad (n > 1)$$

A property of gauge theories is that in four dimensions gauge interactions
are always super-renormalizable
from the weighted power-counting viewpoint

Indeed, for

$$\bar{d} = 1 + \frac{3}{n}$$

the weight of the gauge coupling is

$$[g] = \frac{1}{2} - \frac{1}{2n} > 0 \quad (n > 1)$$

Some fields may have vanishing or negative weights

To have better control over the diagrammatic structure in super-renormalizable theories or theories containing super-renormalizable gauge interactions it is convenient to parametrize the gauge lagrangian as

$$\mathcal{L} = \frac{1}{\alpha} \mathcal{L}_g(gA, g\bar{C}, gC, \lambda_g)$$

To have better control over the diagrammatic structure in super-renormalizable theories or theories containing super-renormalizable gauge interactions it is convenient to parametrize the gauge lagrangian as

$$\mathcal{L} = \frac{1}{\alpha} \mathcal{L}_g(gA, g\bar{C}, gC, \lambda_g)$$

In this way in Feynman diagrams every external leg has a factor g attached to it, which lowers the weight

To have better control over the diagrammatic structure in **super-renormalizable** theories or theories containing **super-renormalizable gauge interactions** it is convenient to parametrize the gauge lagrangian as

$$\mathcal{L} = \frac{1}{\alpha} \mathcal{L}_g(gA, g\bar{C}, gC, \lambda_g)$$

In this way in Feynman diagrams every external leg has **a factor g attached to it**, which **lowers the weight**

Similarly, we can attach possibly weightful coupling constants to scalar fields and fermions as in

$$\begin{aligned} \mathcal{L} = & \frac{1}{\bar{\alpha}_1} \mathcal{L}_1(\bar{g}_1 A) + \frac{1}{\bar{\alpha}_2} \mathcal{L}_2(\bar{g}_2 \psi) + \frac{1}{\bar{\alpha}_3} \mathcal{L}_3(\bar{g}_3 \varphi) + \frac{1}{\bar{a}_3} \mathcal{L}_{12}(\bar{g}_1 A, \bar{g}_2 \psi) \\ & + \frac{1}{\bar{a}_2} \mathcal{L}_{13}(\bar{g}_1 A, \bar{g}_3 \varphi) + \frac{1}{\bar{a}_1} \mathcal{L}_{23}(\bar{g}_2 \psi, \bar{g}_3 \varphi) + \frac{1}{\bar{\alpha}} \mathcal{L}_{123}(\bar{g}_1 A, \bar{g}_2 \psi, \bar{g}_3 \varphi) \end{aligned}$$

To have better control over the diagrammatic structure in **super-renormalizable** theories or theories containing **super-renormalizable gauge interactions** it is convenient to parametrize the gauge lagrangian as

$$\mathcal{L} = \frac{1}{\alpha} \mathcal{L}_g(gA, g\bar{C}, gC, \lambda_g)$$

In this way in Feynman diagrams every external leg has **a factor g attached to it**, which **lowers the weight**

Similarly, we can attach possibly weightful coupling constants to scalar fields and fermions as in

$$\begin{aligned} \mathcal{L} = & \frac{1}{\bar{\alpha}_1} \mathcal{L}_1(\bar{g}_1 A) + \frac{1}{\bar{\alpha}_2} \mathcal{L}_2(\bar{g}_2 \psi) + \frac{1}{\bar{\alpha}_3} \mathcal{L}_3(\bar{g}_3 \varphi) + \frac{1}{\bar{a}_3} \mathcal{L}_{12}(\bar{g}_1 A, \bar{g}_2 \psi) \\ & + \frac{1}{\bar{a}_2} \mathcal{L}_{13}(\bar{g}_1 A, \bar{g}_3 \varphi) + \frac{1}{\bar{a}_1} \mathcal{L}_{23}(\bar{g}_2 \psi, \bar{g}_3 \varphi) + \frac{1}{\bar{\alpha}} \mathcal{L}_{123}(\bar{g}_1 A, \bar{g}_2 \psi, \bar{g}_3 \varphi) \end{aligned}$$

Then counterterms have the form

$$\bar{\alpha}^{L-1} \Delta_1 \mathcal{L}(\bar{g}_1 A, \bar{g}_2 \psi, \bar{g}_3 \varphi)$$

Compatibility with the covariant structure demands

$$[g] \geq [\bar{g}_1], \quad [g\bar{g}_1] \geq [\bar{g}_2^2], \quad [g\bar{g}_1] \geq [\bar{g}_3^2].$$

Compatibility with the **covariant structure** demands

$$[g] \geq [\bar{g}_1], \quad [g\bar{g}_1] \geq [\bar{g}_2^2], \quad [g\bar{g}_1] \geq [\bar{g}_3^2].$$

Polynomiality demands that the weights of

$$\bar{g}_1 F, \bar{g}_2 \psi, \bar{g}_3 \varphi$$

be strictly positive

Compatibility with the covariant structure demands

$$[g] \geq [\bar{g}_1], \quad [g\bar{g}_1] \geq [\bar{g}_2^2], \quad [g\bar{g}_1] \geq [\bar{g}_3^2].$$

Polynomiality demands that the weights of

$$\bar{g}_1 F, \bar{g}_2 \psi, \bar{g}_3 \varphi$$

be strictly positive

Absence of IR divergences in Feynman diagrams at non-exceptional external momenta demands

$$d \geq 4$$

The ones just given are the restrictions to have a **consistent model**

The ones just given are the restrictions to have a **consistent model**

Now we can impose restrictions to have an **interesting model**, namely a model that can explain something the ordinary Standard Model **cannot explain**

The ones just given are the restrictions to have a **consistent model**

Now we can impose restrictions to have an **interesting model**, namely a model that can explain something the ordinary Standard Model **cannot explain**

A **two scalar-two fermion vertex** $(\bar{g}_2^2 \bar{g}_3^2 / \bar{a}_1) \varphi^2 \bar{\psi} \psi$ is renormalizable if

$$[g_{2,3}] \leq 1 - \frac{3}{2n}$$

The ones just given are the restrictions to have a **consistent model**

Now we can impose restrictions to have an **interesting model**, namely a model that can explain something the ordinary Standard Model **cannot explain**

A **two scalar-two fermion vertex** $(\bar{g}_2^2 \bar{g}_3^2 / \bar{a}_1) \varphi^2 \bar{\psi} \psi$ is renormalizable if

$$[g_{2,3}] \leq 1 - \frac{3}{2n}$$

A **four-fermion vertex** $\bar{\psi} \psi \bar{\psi} \psi$ is renormalizable if

$$[g_2] \leq \frac{1}{2} - \frac{3}{2n}$$

We can also make the simplifying assumption

$$g_1 = g$$

We can also make the simplifying assumption

$$g_1 = g$$

The combination of **all requirements** listed so far gives, in four dimensions

$$\frac{1}{2} - \frac{3}{2n} < [g_3] \leq \frac{1}{2} - \frac{1}{2n}$$
$$0 \leq [g_2] \leq \frac{1}{2} - \frac{3}{2n}$$

We can also make the simplifying assumption

$$g_1 = g$$

The combination of **all requirements** listed so far gives, in four dimensions

$$\frac{1}{2} - \frac{3}{2n} < [g_3] \leq \frac{1}{2} - \frac{1}{2n}$$
$$0 \leq [g_2] \leq \frac{1}{2} - \frac{3}{2n}$$

Solutions exist **for every** $n \geq 3$

We can also make the simplifying assumption

$$g_1 = g$$

The combination of **all requirements** listed so far gives, in four dimensions

$$\frac{1}{2} - \frac{3}{2n} < [g_3] \leq \frac{1}{2} - \frac{1}{2n}$$
$$0 \leq [g_2] \leq \frac{1}{2} - \frac{3}{2n}$$

Solutions exist **for every** $n \geq 3$

$n = \text{odd}$ is necessary to describe **chiral fermions**

We can also make the simplifying assumption

$$g_1 = g$$

The combination of **all requirements** listed so far gives, in four dimensions

$$\frac{1}{2} - \frac{3}{2n} < [g_3] \leq \frac{1}{2} - \frac{1}{2n}$$
$$0 \leq [g_2] \leq \frac{1}{2} - \frac{3}{2n}$$

Solutions exist **for every** $n \geq 3$

$n = \text{odd}$ is necessary to describe **chiral fermions**

The simplest choice is

$$\bar{d} = 2, \quad n = 3, \quad [g] = [g_1] = [g_3] = \frac{1}{3}, \quad [g_2] = 0.$$

The case $\mathfrak{d} = 2$, $n = 3$, $[g] = [g_1] = [g_3] = \frac{1}{3}$, $[g_2] = 0$.

The case $\bar{d} = 2$, $n = 3$, $[g] = [g_1] = [g_3] = \frac{1}{3}$, $[g_2] = 0$.

allows us to formulate a consistent Lorentz violating extended Standard Model that contains both the dimension-5 vertex

$$\mathcal{L}_{LH} = \frac{\bar{g}^2}{4\Lambda_L} (LH)^2$$

$$(LH)^2 \equiv \sum_{a,b=1}^3 Y_{ab} \varepsilon_{ij} L_i^{\alpha a} H_j \varepsilon_{\alpha\beta} \varepsilon_{kl} L_k^{\beta b} H_l + \text{h.c.}$$

that gives **Majorana masses** to **left-handed** neutrinos after symmetry breaking,

The case $\bar{d} = 2$, $n = 3$, $[g] = [g_1] = [g_3] = \frac{1}{3}$, $[g_2] = 0$.

allows us to formulate a consistent Lorentz violating extended Standard Model that contains both the dimension-5 vertex

$$\mathcal{L}_{LH} = \frac{\bar{g}^2}{4\Lambda_L} (LH)^2$$

$$(LH)^2 \equiv \sum_{a,b=1}^3 Y_{ab} \varepsilon_{ij} L_i^{\alpha a} H_j \varepsilon_{\alpha\beta} \varepsilon_{kl} L_k^{\beta b} H_l + \text{h.c.}$$

that gives **Majorana masses** to **left-handed** neutrinos after symmetry breaking, and the four fermion interactions

$$\mathcal{L}_{4f} \sim \frac{Y_f}{\Lambda_L^2} \bar{\psi}\psi\bar{\psi}\psi$$

that can describe proton decay.

The case $\bar{d} = 2$, $n = 3$, $[g] = [g_1] = [g_3] = \frac{1}{3}$, $[g_2] = 0$.

allows us to formulate a consistent Lorentz violating extended Standard Model that contains both the dimension-5 vertex

$$\mathcal{L}_{LH} = \frac{\bar{g}^2}{4\Lambda_L} (LH)^2$$

$$(LH)^2 \equiv \sum_{a,b=1}^3 Y_{ab} \varepsilon_{ij} L_i^{\alpha a} H_j \varepsilon_{\alpha\beta} \varepsilon_{kl} L_k^{\beta b} H_l + \text{h.c.}$$

that gives **Majorana masses** to **left-handed** neutrinos after symmetry breaking, and the four fermion interactions

$$\mathcal{L}_{4f} \sim \frac{Y_f}{\Lambda_L^2} \bar{\psi}\psi\bar{\psi}\psi$$

that can describe proton decay.

Such vertices **are renormalizable by weighted power counting**

The case $\bar{d} = 2$, $n = 3$, $[g] = [g_1] = [g_3] = \frac{1}{3}$, $[g_2] = 0$.

allows us to formulate a consistent Lorentz violating extended Standard Model that contains both the dimension-5 vertex

$$\mathcal{L}_{LH} = \frac{\bar{g}^2}{4\Lambda_L} (LH)^2$$

$$(LH)^2 \equiv \sum_{a,b=1}^3 Y_{ab} \varepsilon_{ij} L_i^{\alpha a} H_j \varepsilon_{\alpha\beta} \varepsilon_{kl} L_k^{\beta b} H_l + \text{h.c.}$$

that gives **Majorana masses** to **left-handed** neutrinos after symmetry breaking, and the four fermion interactions

$$\mathcal{L}_{4f} \sim \frac{Y_f}{\Lambda_L^2} \bar{\psi}\psi\bar{\psi}\psi$$

that can describe proton decay.

Such vertices **are renormalizable by weighted power counting**

Matching the vertex $(LH)^2$ with estimates of the electron neutrino Majorana mass the **scale of Lorentzn violation** has roughly the value

$$\Lambda_L \sim 10^{14} \text{GeV}$$

The (simplified) model reads

$$\begin{aligned}\mathcal{L}' = & \mathcal{L}'_Q + \mathcal{L}_{\text{kin}f} + \mathcal{L}'_H + \mathcal{L}_Y - \frac{\bar{g}^2}{4\Lambda_L}(LH)^2 - \sum_{I=1}^5 \frac{1}{\Lambda_L^2} g \bar{D} \bar{F} (\bar{\chi}_I \bar{\gamma} \chi_I) + \frac{Y_f}{\Lambda_L^2} \bar{\psi} \psi \bar{\psi} \psi - \frac{g}{\Lambda_L^2} \bar{F}^3 \\ & - \frac{1}{\Lambda_L^2} g \bar{g} \bar{\psi} \psi \bar{F} H - \frac{1}{\Lambda_L^2} (\bar{g}^3 \bar{\psi} \psi H^3 + \bar{g}^2 \bar{\psi} \bar{D} \psi H^2 + \bar{g} \bar{\psi} \bar{D}^2 \psi H) - \frac{1}{\Lambda_L^4} (g \bar{D}^2 \bar{F} + g^2 \bar{F}^2) H^\dagger H,\end{aligned}$$

The (simplified) model reads

$$\begin{aligned} \mathcal{L}' = & \mathcal{L}'_Q + \mathcal{L}_{\text{kin}f} + \mathcal{L}'_H + \mathcal{L}_Y - \frac{\bar{g}^2}{4\Lambda_L} (LH)^2 - \sum_{I=1}^5 \frac{1}{\Lambda_L^2} g \bar{D} \bar{F} (\bar{\chi}_I \bar{\gamma} \chi_I) + \frac{Y_f}{\Lambda_L^2} \bar{\psi} \psi \bar{\psi} \psi - \frac{g}{\Lambda_L^2} \bar{F}^3 \\ & - \frac{1}{\Lambda_L^2} g \bar{g} \bar{\psi} \psi \bar{F} H - \frac{1}{\Lambda_L^2} (\bar{g}^3 \bar{\psi} \psi H^3 + \bar{g}^2 \bar{\psi} \bar{D} \psi H^2 + \bar{g} \bar{\psi} \bar{D}^2 \psi H) - \frac{1}{\Lambda_L^4} (g \bar{D}^2 \bar{F} + g^2 \bar{F}^2) H^\dagger H, \end{aligned}$$

where

$$\mathcal{L}'_Q = \frac{1}{4} \sum_G (2F_{\hat{\mu}\bar{\nu}}^G F_{\hat{\mu}\bar{\nu}}^G - F_{\bar{\mu}\bar{\nu}}^G \tau'^G(\bar{\Upsilon}) F_{\bar{\mu}\bar{\nu}}^G),$$

The (simplified) model reads

$$\begin{aligned} \mathcal{L}' = & \mathcal{L}'_Q + \mathcal{L}_{\text{kin}f} + \mathcal{L}'_H + \mathcal{L}_Y - \frac{\bar{g}^2}{4\Lambda_L} (LH)^2 - \sum_{I=1}^5 \frac{1}{\Lambda_L^2} g \bar{D} \bar{F} (\bar{\chi}_I \bar{\gamma} \chi_I) + \frac{Y_f}{\Lambda_L^2} \bar{\psi} \psi \bar{\psi} \psi - \frac{g}{\Lambda_L^2} \bar{F}^3 \\ & - \frac{1}{\Lambda_L^2} g \bar{g} \bar{\psi} \psi \bar{F} H - \frac{1}{\Lambda_L^2} (\bar{g}^3 \bar{\psi} \psi H^3 + \bar{g}^2 \bar{\psi} \bar{D} \psi H^2 + \bar{g} \bar{\psi} \bar{D}^2 \psi H) - \frac{1}{\Lambda_L^4} (g \bar{D}^2 \bar{F} + g^2 \bar{F}^2) H^\dagger H, \end{aligned}$$

where

$$\mathcal{L}'_Q = \frac{1}{4} \sum_G (2F_{\hat{\mu}\bar{\nu}}^G F_{\hat{\mu}\bar{\nu}}^G - F_{\bar{\mu}\bar{\nu}}^G \tau'^G(\bar{\Upsilon}) F_{\bar{\mu}\bar{\nu}}^G),$$

$$\mathcal{L}'_H = \mathcal{L}_H - \frac{\lambda_4^{(3)} \bar{g}^2}{4\Lambda_L^2} |H|^2 |\bar{D}_{\bar{\mu}} H|^2 - \frac{\lambda_4^{(2)} \bar{g}^2}{4\Lambda_L^2} |H^\dagger \bar{D}_{\bar{\mu}} H|^2 - \frac{\bar{g}^2}{4\Lambda_L^2} \left[\lambda_4^{(1)} (H^\dagger \bar{D}_{\bar{\mu}} H)^2 + \text{h.c.} \right] - \frac{\lambda_6 \bar{g}^4}{36\Lambda_L^2} |H|^6$$

The (simplified) model reads

$$\begin{aligned}\mathcal{L}' = & \mathcal{L}'_Q + \mathcal{L}_{\text{kin}f} + \mathcal{L}'_H + \mathcal{L}_Y - \frac{\bar{g}^2}{4\Lambda_L}(LH)^2 - \sum_{I=1}^5 \frac{1}{\Lambda_L^2} g \bar{D} \bar{F} (\bar{\chi}_I \bar{\gamma} \chi_I) + \frac{Y_f}{\Lambda_L^2} \bar{\psi} \psi \bar{\psi} \psi - \frac{g}{\Lambda_L^2} \bar{F}^3 \\ & - \frac{1}{\Lambda_L^2} g \bar{g} \bar{\psi} \psi \bar{F} H - \frac{1}{\Lambda_L^2} (\bar{g}^3 \bar{\psi} \psi H^3 + \bar{g}^2 \bar{\psi} \bar{D} \psi H^2 + \bar{g} \bar{\psi} \bar{D}^2 \psi H) - \frac{1}{\Lambda_L^4} (g \bar{D}^2 \bar{F} + g^2 \bar{F}^2) H^\dagger H,\end{aligned}$$

where

$$\mathcal{L}'_Q = \frac{1}{4} \sum_G (2F_{\hat{\mu}\bar{\nu}}^G F_{\hat{\mu}\bar{\nu}}^G - F_{\hat{\mu}\bar{\nu}}^G \tau'^G(\bar{\Upsilon}) F_{\hat{\mu}\bar{\nu}}^G),$$

$$\mathcal{L}'_H = \mathcal{L}_H - \frac{\lambda_4^{(3)} \bar{g}^2}{4\Lambda_L^2} |H|^2 |\bar{D}_{\bar{\mu}} H|^2 - \frac{\lambda_4^{(2)} \bar{g}^2}{4\Lambda_L^2} |H^\dagger \bar{D}_{\bar{\mu}} H|^2 - \frac{\bar{g}^2}{4\Lambda_L^2} \left[\lambda_4^{(1)} (H^\dagger \bar{D}_{\bar{\mu}} H)^2 + \text{h.c.} \right] - \frac{\lambda_6 \bar{g}^4}{36\Lambda_L^2} |H|^6$$

$$\mathcal{L}_{\text{kin}f} = \sum_{a,b=1}^3 \sum_{I=1}^5 \bar{\chi}_I^a i \left(\delta^{ab} \hat{\not{D}} - \frac{b_0^{Iab}}{\Lambda_L^2} \bar{\not{D}}^3 + b_1^{Iab} \bar{\not{D}} \right) \chi_I^b,$$

The (simplified) model reads

$$\begin{aligned}\mathcal{L}' = & \mathcal{L}'_Q + \mathcal{L}_{\text{kin}f} + \mathcal{L}'_H + \mathcal{L}_Y - \frac{\bar{g}^2}{4\Lambda_L}(LH)^2 - \sum_{I=1}^5 \frac{1}{\Lambda_L^2} g \bar{D} \bar{F} (\bar{\chi}_I \bar{\gamma} \chi_I) + \frac{Y_f}{\Lambda_L^2} \bar{\psi} \psi \bar{\psi} \psi - \frac{g}{\Lambda_L^2} \bar{F}^3 \\ & - \frac{1}{\Lambda_L^2} g \bar{g} \bar{\psi} \psi \bar{F} H - \frac{1}{\Lambda_L^2} (\bar{g}^3 \bar{\psi} \psi H^3 + \bar{g}^2 \bar{\psi} \bar{D} \psi H^2 + \bar{g} \bar{\psi} \bar{D}^2 \psi H) - \frac{1}{\Lambda_L^4} (g \bar{D}^2 \bar{F} + g^2 \bar{F}^2) H^\dagger H,\end{aligned}$$

where

$$\mathcal{L}'_Q = \frac{1}{4} \sum_G (2F_{\hat{\mu}\bar{\nu}}^G F_{\hat{\mu}\bar{\nu}}^G - F_{\hat{\mu}\bar{\nu}}^G \tau'^G(\bar{\Upsilon}) F_{\hat{\mu}\bar{\nu}}^G),$$

$$\mathcal{L}'_H = \mathcal{L}_H - \frac{\lambda_4^{(3)} \bar{g}^2}{4\Lambda_L^2} |H|^2 |\bar{D}_{\bar{\mu}} H|^2 - \frac{\lambda_4^{(2)} \bar{g}^2}{4\Lambda_L^2} |H^\dagger \bar{D}_{\bar{\mu}} H|^2 - \frac{\bar{g}^2}{4\Lambda_L^2} \left[\lambda_4^{(1)} (H^\dagger \bar{D}_{\bar{\mu}} H)^2 + \text{h.c.} \right] - \frac{\lambda_6 \bar{g}^4}{36\Lambda_L^2} |H|^6$$

$$\mathcal{L}_{\text{kin}f} = \sum_{a,b=1}^3 \sum_{I=1}^5 \bar{\chi}_I^a i \left(\delta^{ab} \hat{\mathcal{P}} - \frac{b_0^{Iab}}{\Lambda_L^2} \bar{\mathcal{P}}^3 + b_1^{Iab} \bar{\mathcal{P}} \right) \chi_I^b,$$

$$\mathcal{L}_Y = -\bar{g} \Omega_i H^i + \text{h.c.}, \quad \Omega_i = \sum_{a,b=1}^3 Y_1^{ab} \bar{L}^{ai} \ell_R^b + Y_2^{ab} \bar{u}_R^a Q_L^{bj} \varepsilon^{ji} + Y_3^{ab} \bar{Q}_L^{ai} d_R^b,$$

At low energies we have the Colladay-Kostelecky **Standard-Model Extension**

The (simplified) model reads

$$\mathcal{L}' = \mathcal{L}'_Q + \mathcal{L}_{\text{kinf}} + \mathcal{L}'_H + \mathcal{L}_Y - \frac{\bar{g}^2}{4\Lambda_L}(LH)^2 - \sum_{I=1}^5 \frac{1}{\Lambda_L^2} g \bar{D} \bar{F} (\bar{\chi}_I \bar{\gamma} \chi_I) + \frac{Y_f}{\Lambda_L^2} \bar{\psi} \psi \bar{\psi} \psi - \frac{g}{\Lambda_L^2} \bar{F}^3$$

$$- \frac{1}{\Lambda_L^2} g \bar{g} \bar{\psi} \psi \bar{F} H - \frac{1}{\Lambda_L^2} (\bar{g}^3 \bar{\psi} \psi H^3 + \bar{g}^2 \bar{\psi} \bar{D} \psi H^2 + \bar{g} \bar{\psi} \bar{D}^2 \psi H) - \frac{1}{\Lambda_L^4} (g \bar{D}^2 \bar{F} + g^2 \bar{F}^2) H^\dagger H,$$

where

$$\mathcal{L}'_Q = \frac{1}{4} \sum_G (2F_{\hat{\mu}\bar{\nu}}^G F_{\hat{\mu}\bar{\nu}}^G - F_{\hat{\mu}\bar{\nu}}^G \tau'^G(\bar{\Upsilon}) F_{\hat{\mu}\bar{\nu}}^G),$$

$$\mathcal{L}'_H = \mathcal{L}_H - \frac{\lambda_4^{(3)} \bar{g}^2}{4\Lambda_L^2} |H|^2 |\bar{D}_{\bar{\mu}} H|^2 - \frac{\lambda_4^{(2)} \bar{g}^2}{4\Lambda_L^2} |H^\dagger \bar{D}_{\bar{\mu}} H|^2 - \frac{\bar{g}^2}{4\Lambda_L^2} [\lambda_4^{(1)} (H^\dagger \bar{D}_{\bar{\mu}} H)^2 + \text{h.c.}] - \frac{\lambda_6 \bar{g}^4}{36\Lambda_L^2} |H|^6$$

$$\mathcal{L}_{\text{kinf}} = \sum_{a,b=1}^3 \sum_{I=1}^5 \bar{\chi}_I^a i \left(\delta^{ab} \hat{\mathcal{P}} - \frac{b_0^{Iab}}{\Lambda_L^2} \bar{\mathcal{P}}^3 + b_1^{Iab} \bar{\mathcal{P}} \right) \chi_I^b,$$

$$\mathcal{L}_Y = -\bar{g} \Omega_i H^i + \text{h.c.}, \quad \Omega_i = \sum_{a,b=1}^3 Y_1^{ab} \bar{L}^{ai} \ell_R^b + Y_2^{ab} \bar{u}_R^a Q_L^{bj} \varepsilon^{ji} + Y_3^{ab} \bar{Q}_L^{ai} d_R^b,$$

At low energies we have the Colladay-Kostelecky **Standard-Model Extension**

It can be shown that the **gauge anomalies vanish**, since they coincide with those of the Standard Model

Low-energy Lorentz recovery

Low-energy Lorentz recovery

Once Lorentz invariance is broken at **some energy**, it is not automatic to **recover** **it** at **lower energies**, because renormalization makes couplings run independently

Low-energy Lorentz recovery

Once Lorentz invariance is broken at **some energy**, it is not automatic to **recover it** at **lower energies**, because renormalization makes couplings run independently

For example, we can find at low energies

$$\bar{\psi} \gamma^\mu (\hat{\partial}_\mu + \eta \bar{\partial}_\mu) \psi$$

Low-energy Lorentz recovery

Once Lorentz invariance is broken at **some energy**, it is not automatic to **recover** it at **lower energies**, because renormalization makes couplings run independently

For example, we can find at low energies

$$\bar{\psi} \gamma^\mu (\hat{\partial}_\mu + \eta \bar{\partial}_\mu) \psi$$

The **Lorentz invariant surface** is **RG invariant**. Nielsen conjectured long ago that it could also be **IR attractive** (evidence is not strong)

Low-energy Lorentz recovery

Once Lorentz invariance is broken at **some energy**, it is not automatic to **recover** it at **lower energies**, because renormalization makes couplings run independently

For example, we can find at low energies

$$\bar{\psi} \gamma^\mu (\hat{\partial}_\mu + \eta \bar{\partial}_\mu) \psi$$

The **Lorentz invariant surface** is **RG invariant**. Nielsen conjectured long ago that it could also be **IR attractive** (evidence is not strong)

In general, we must argue as in the case of asymptotic safety: all low-energy

Lorentz violating parameters that **flow to** the surface can be kept, all those that **flow away** from the surface must be set to zero (**fine tuning**)

Given that the model contains four fermion interactions at the fundamental level, we may ask if it is possible to describe the known low-energy physics without introducing elementary scalars, in the Nambu—Jona-Lasinio spirit.

Given that the model contains four fermion interactions at the fundamental level, we may ask if it is possible to describe the known low-energy physics without introducing elementary scalars, in the Nambu—Jona-Lasinio spirit.

Then the Higgs field must be composite and arise as a low-energy effect of the four fermion vertices.

Given that the model contains four fermion interactions at the fundamental level, we may ask if it is possible to describe the known low-energy physics without introducing elementary scalars, in the Nambu—Jona-Lasinio spirit.

Then the Higgs field must be composite and arise as a low-energy effect of the four fermion vertices.

Suppressing the elementary Higgs field, we obtain the scalarless model

$$\mathcal{L}_{\text{noH}} = \mathcal{L}'_Q + \mathcal{L}_{\text{kinf}} - \sum_{I=1}^5 \frac{1}{\Lambda_L^2} g \bar{D} \bar{F} (\bar{\chi}_I \bar{\gamma} \chi_I) + \frac{Y_f}{\Lambda_L^2} \bar{\psi} \psi \bar{\psi} \psi - \frac{g}{\Lambda_L^2} \bar{F}^3,$$

Given that the model contains four fermion interactions at the fundamental level, we may ask if it is possible to describe the known low-energy physics without introducing elementary scalars, in the Nambu—Jona-Lasinio spirit.

Then the Higgs field must be composite and arise as a low-energy effect of the four fermion vertices.

Suppressing the elementary Higgs field, we obtain the scalarless model

$$\mathcal{L}_{\text{noH}} = \mathcal{L}'_Q + \mathcal{L}_{\text{kinf}} - \sum_{I=1}^5 \frac{1}{\Lambda_L^2} g \bar{D} \bar{F} (\bar{\chi}_I \bar{\gamma} \chi_I) + \frac{Y_f}{\Lambda_L^2} \bar{\psi} \psi \bar{\psi} \psi - \frac{g}{\Lambda_L^2} \bar{F}^3,$$

Moreover, since the gauge interactions are super-renormalizable, in the weighted power counting framework, at very high energies they become negligible and the model reduces to a four fermion model in two weighted dimensions, described by the lagrangian

$$\mathcal{L}_{4f} = \sum_{a,b=1}^3 \sum_{I=1}^5 \bar{\chi}_I^a i \left(\delta^{ab} \hat{\partial} - \frac{b_0^{Iab}}{\Lambda_L^2} \bar{\partial}^3 + b_1^{Iab} \bar{\partial} \right) \chi_I^b + \frac{Y_f}{\Lambda_L^2} \bar{\psi} \psi \bar{\psi} \psi.$$

Consider the t - b model

$$\mathcal{L}_q = \sum_{I=1}^{N_c} \bar{\psi}_I \left(\Gamma^\mu i \left(\hat{\partial}_\mu + \bar{\partial}_\mu - \bar{\partial}_\mu \frac{\bar{\partial}^2}{\Lambda_L^2} \right) - M \right) \psi_I - V_2(M),$$

Consider the t - b model

$$\mathcal{L}_q = \sum_{I=1}^{N_c} \bar{\psi}_I \left(\Gamma^\mu i \left(\hat{\partial}_\mu + \bar{\partial}_\mu - \bar{\partial}_\mu \frac{\bar{\partial}^2}{\Lambda_L^2} \right) - M \right) \psi_I - V_2(M),$$

in the large N_c limit, where

$$M = \begin{pmatrix} \tau & \rho_R \\ \rho_L & \tau^\dagger \end{pmatrix}, \quad \psi = \begin{pmatrix} Q_L^i \\ Q_R^k \end{pmatrix}$$

$$\Lambda_L^{-2} V_2(M) = \text{tr}[\tau \tau^\dagger C] + \frac{1}{2g_L^2} \text{tr}[\rho_L^2] + \frac{1}{2g_L'^2} (\text{tr}[\rho_L])^2 + g_R^{kl} \text{tr}[\rho_L] \rho_R^{kl} + \frac{1}{2} g_{RR}^{klmn} \rho_R^{kl} \rho_R^{mn}$$

Consider the t - b model

$$\mathcal{L}_q = \sum_{I=1}^{N_c} \bar{\psi}_I \left(\Gamma^\mu i \left(\hat{\partial}_\mu + \bar{\partial}_\mu - \bar{\partial}_\mu \frac{\bar{\partial}^2}{\Lambda_L^2} \right) - M \right) \psi_I - V_2(M),$$

In the large N_c limit, where

$$M = \begin{pmatrix} \tau & \rho_R \\ \rho_L & \tau^\dagger \end{pmatrix}, \quad \psi = \begin{pmatrix} Q_L^i \\ Q_R^k \end{pmatrix}$$

$$\Lambda_L^{-2} V_2(M) = \text{tr}[\tau \tau^\dagger C] + \frac{1}{2g_L^2} \text{tr}[\rho_L^2] + \frac{1}{2g_L'^2} (\text{tr}[\rho_L])^2 + g_R^{kl} \text{tr}[\rho_L] \rho_R^{kl} + \frac{1}{2} g_{RR}^{klmn} \rho_R^{kl} \rho_R^{mn}$$

We can prove that

$$\tau = \begin{pmatrix} m_t & 0 \\ 0 & m_b \end{pmatrix} \equiv \tau_0, \quad \rho_L = \rho_R = 0.$$

is a minimum of the effective potential

Consider the t - b model

$$\mathcal{L}_q = \sum_{I=1}^{N_c} \bar{\psi}_I \left(\Gamma^\mu i \left(\hat{\partial}_\mu + \bar{\partial}_\mu - \bar{\partial}_\mu \frac{\bar{\partial}^2}{\Lambda_L^2} \right) - M \right) \psi_I - V_2(M),$$

In the large N_c limit, where

$$M = \begin{pmatrix} \tau & \rho_R \\ \rho_L & \tau^\dagger \end{pmatrix}, \quad \psi = \begin{pmatrix} Q_L^i \\ Q_R^k \end{pmatrix}$$

$$\Lambda_L^{-2} V_2(M) = \text{tr}[\tau \tau^\dagger C] + \frac{1}{2g_L^2} \text{tr}[\rho_L^2] + \frac{1}{2g_L'^2} (\text{tr}[\rho_L])^2 + g_R^{kl} \text{tr}[\rho_L] \rho_R^{kl} + \frac{1}{2} g_{RR}^{klmn} \rho_R^{kl} \rho_R^{mn}$$

We can prove that

$$\tau = \begin{pmatrix} m_t & 0 \\ 0 & m_b \end{pmatrix} \equiv \tau_0, \quad \rho_L = \rho_R = 0.$$

is a minimum of the effective potential and gives masses to the fermions

The Lorentz violating fermionic mass terms are not turned on, so the Lorentz violation remains highly suppressed.

The Lorentz violating fermionic mass terms are not turned on, so the Lorentz violation remains highly suppressed.

Moreover, we can read the bound states from the effective potential. We find

$$\Gamma_{\eta\eta} = N_c \sum_{i,j=t,b} \{ 2\eta_{ij}(p)\bar{\eta}_{ij}(-p)(p^2 f'_{ij} - m_j^2 f_{ij}) - m_i m_j f_{ij} [\eta_{ij}(p)\eta_{ji}(-p) + \bar{\eta}_{ij}(p)\bar{\eta}_{ji}(-p)] \}$$

The Lorentz violating fermionic mass terms are not turned on, so the Lorentz violation remains highly suppressed.

Moreover, we can read the bound states from the effective potential. We find

$$\Gamma_{\eta\eta} = N_c \sum_{i,j=t,b} \{ 2\eta_{ij}(p)\bar{\eta}_{ij}(-p)(p^2 f'_{ij} - m_j^2 f_{ij}) - m_i m_j f_{ij} [\eta_{ij}(p)\eta_{ji}(-p) + \bar{\eta}_{ij}(p)\bar{\eta}_{ji}(-p)] \}$$

where

$$\tau = \tau_0 + \eta \quad f_{ii} \sim \frac{1}{(4\pi)^2} \ln \frac{\Lambda_L^2}{m_i^2}, \quad f_{tb} = f_{bt} \sim f_{tt}, \quad f'_{ij} \sim \frac{1}{2} f_{ij}, \quad f''_{ij} \sim \frac{1}{6} f_{ij}$$

The Lorentz violating fermionic mass terms are not turned on, so the Lorentz violation remains highly suppressed.

Moreover, we can read the bound states from the effective potential. We find

$$\Gamma_{\eta\eta} = N_c \sum_{i,j=t,b} \{ 2\eta_{ij}(p)\bar{\eta}_{ij}(-p)(p^2 f'_{ij} - m_j^2 f_{ij}) - m_i m_j f_{ij} [\eta_{ij}(p)\eta_{ji}(-p) + \bar{\eta}_{ij}(p)\bar{\eta}_{ji}(-p)] \}$$

where

$$\tau = \tau_0 + \eta \quad f_{ii} \sim \frac{1}{(4\pi)^2} \ln \frac{\Lambda_L^2}{m_i^2}, \quad f_{tb} = f_{bt} \sim f_{tt}, \quad f'_{ij} \sim \frac{1}{2} f_{ij}, \quad f''_{ij} \sim \frac{1}{6} f_{ij}$$

Studying the poles we find

1) two neutral massive bound states of squared masses

$$2m_t^2 \frac{f_{tt}}{f'_{tt}} = 4m_t^2, \quad 2m_b^2 \frac{f_{bb}}{f'_{bb}} = 4m_b^2;$$

The Lorentz violating fermionic mass terms are not turned on, so the Lorentz violation remains highly suppressed.

Moreover, we can read the bound states from the effective potential. We find

$$\Gamma_{\eta\eta} = N_c \sum_{i,j=t,b} \{ 2\eta_{ij}(p)\bar{\eta}_{ij}(-p)(p^2 f'_{ij} - m_j^2 f_{ij}) - m_i m_j f_{ij} [\eta_{ij}(p)\eta_{ji}(-p) + \bar{\eta}_{ij}(p)\bar{\eta}_{ji}(-p)] \}$$

where

$$\tau = \tau_0 + \eta \quad f_{ii} \sim \frac{1}{(4\pi)^2} \ln \frac{\Lambda_L^2}{m_i^2}, \quad f_{tb} = f_{bt} \sim f_{tt}, \quad f'_{ij} \sim \frac{1}{2} f_{ij}, \quad f''_{ij} \sim \frac{1}{6} f_{ij}$$

Studying the poles we find

1) two neutral massive bound states of squared masses

$$2m_t^2 \frac{f_{tt}}{f'_{tt}} = 4m_t^2, \quad 2m_b^2 \frac{f_{bb}}{f'_{bb}} = 4m_b^2;$$

2) two neutral and two charged massless states, which are the Goldstone bosons;

The Lorentz violating fermionic mass terms are not turned on, so the Lorentz violation remains highly suppressed.

Moreover, we can read the bound states from the effective potential. We find

$$\Gamma_{\eta\eta} = N_c \sum_{i,j=t,b} \{ 2\eta_{ij}(p)\bar{\eta}_{ij}(-p)(p^2 f'_{ij} - m_j^2 f_{ij}) - m_i m_j f_{ij} [\eta_{ij}(p)\eta_{ji}(-p) + \bar{\eta}_{ij}(p)\bar{\eta}_{ji}(-p)] \}$$

where

$$\tau = \tau_0 + \eta \quad f_{ii} \sim \frac{1}{(4\pi)^2} \ln \frac{\Lambda_L^2}{m_i^2}, \quad f_{tb} = f_{bt} \sim f_{tt}, \quad f'_{ij} \sim \frac{1}{2} f_{ij}, \quad f''_{ij} \sim \frac{1}{6} f_{ij}$$

Studying the poles we find

1) two neutral massive bound states of squared masses

$$2m_t^2 \frac{f_{tt}}{f'_{tt}} = 4m_t^2, \quad 2m_b^2 \frac{f_{bb}}{f'_{bb}} = 4m_b^2;$$

2) two neutral and two charged massless states, which are the Goldstone bosons;

3) two charged massive bound states of squared masses

$$f_{tb} \left(\frac{m_t^2}{f'_{bt}} + \frac{m_b^2}{f'_{tb}} \right) \sim 2m_t^2.$$

When gauge interactions are turned on, the Goldstone bosons are “eaten” by W 's and Z , as usual.

When gauge interactions are turned on, the Goldstone bosons are “eaten” by W’s and Z, as usual. Precisely, the effective potential becomes

$$\Gamma_{\eta\eta} - \Gamma_{\phi\phi} + \frac{N_c}{f_W} (\partial_\mu \phi^+ - g f_W W_\mu^+) (\partial^\mu \phi^- - g f_W W^{\mu-}) + \frac{N_c}{2f_Z} (\partial_\mu \phi^0 - \tilde{g} f_Z Z_\mu) (\partial^\mu \phi^0 - \tilde{g} f_Z Z^\mu)$$

When gauge interactions are turned on, the Goldstone bosons are “eaten” by W’s and Z, as usual. Precisely, the effective potential becomes

$$\Gamma_{\eta\eta} - \Gamma_{\phi\phi} + \frac{N_c}{f_W} (\partial_\mu \phi^+ - g f_W W_\mu^+) (\partial^\mu \phi^- - g f_W W^{\mu-}) + \frac{N_c}{2f_Z} (\partial_\mu \phi^0 - \tilde{g} f_Z Z_\mu) (\partial^\mu \phi^0 - \tilde{g} f_Z Z^\mu)$$

where

$$\phi^+ = i\sqrt{2}(m_t f'_{tb} \eta_{tb} - m_b f'_{bt} \bar{\eta}_{bt}), \quad \phi^0 = \frac{i}{2}(m_t f_{tt}(\eta_{tt} - \bar{\eta}_{tt}) - m_b f_{bb}(\eta_{bb} - \bar{\eta}_{bb})),$$

$$f_W = m_t^2 f'_{tb} + m_b^2 f'_{bt}, \quad f_Z = \frac{1}{2}(m_t^2 f_{tt} + m_b^2 f_{bb}).$$

When gauge interactions are turned on, the Goldstone bosons are “eaten” by W’s and Z, as usual. Precisely, the effective potential becomes

$$\Gamma_{\eta\eta} - \Gamma_{\phi\phi} + \frac{N_c}{f_W}(\partial_\mu\phi^+ - gf_W W_\mu^+)(\partial^\mu\phi^- - gf_W W^{\mu-}) + \frac{N_c}{2f_Z}(\partial_\mu\phi^0 - \tilde{g}f_Z Z_\mu)(\partial^\mu\phi^0 - \tilde{g}f_Z Z^\mu)$$

where

$$\phi^+ = i\sqrt{2}(m_t f'_{tb}\eta_{tb} - m_b f'_{bt}\bar{\eta}_{bt}), \quad \phi^0 = \frac{i}{2}(m_t f_{tt}(\eta_{tt} - \bar{\eta}_{tt}) - m_b f_{bb}(\eta_{bb} - \bar{\eta}_{bb})),$$

$$f_W = m_t^2 f'_{tb} + m_b^2 f'_{bt}, \quad f_Z = \frac{1}{2}(m_t^2 f_{tt} + m_b^2 f_{bb}).$$

Choosing the unitary gauge fixing we find the gauge-boson mass terms.

$$\phi^\pm = \phi^0 = 0$$

When gauge interactions are turned on, the Goldstone bosons are “eaten” by W’s and Z, as usual. Precisely, the effective potential becomes

$$\Gamma_{\eta\eta} - \Gamma_{\phi\phi} + \frac{N_c}{f_W} (\partial_\mu \phi^+ - g f_W W_\mu^+) (\partial^\mu \phi^- - g f_W W^{\mu-}) + \frac{N_c}{2f_Z} (\partial_\mu \phi^0 - \tilde{g} f_Z Z_\mu) (\partial^\mu \phi^0 - \tilde{g} f_Z Z^\mu)$$

where

$$\phi^+ = i\sqrt{2}(m_t f'_{tb} \eta_{tb} - m_b f'_{bt} \bar{\eta}_{bt}), \quad \phi^0 = \frac{i}{2}(m_t f_{tt}(\eta_{tt} - \bar{\eta}_{tt}) - m_b f_{bb}(\eta_{bb} - \bar{\eta}_{bb})),$$

$$f_W = m_t^2 f'_{tb} + m_b^2 f'_{bt}, \quad f_Z = \frac{1}{2}(m_t^2 f_{tt} + m_b^2 f_{bb}).$$

Choosing the unitary gauge fixing $\phi^\pm = \phi^0 = 0$
we find the gauge-boson mass terms. The masses are

$$m_W^2 = N_c g^2 f_W \sim \frac{N_c g^2}{32\pi^2} m_t^2 \ln \frac{\Lambda_L^2}{m_t^2}, \quad m_Z^2 = N_c \tilde{g}^2 f_Z \sim \frac{\tilde{g}^2}{g^2} m_W^2.$$

In particular, such relations give

$$\frac{1}{G_F} = \frac{N_c m_t^2}{4\pi^2 \sqrt{2}} \ln \frac{\Lambda_L^2}{m_t^2}.$$

In particular, such relations give

$$\frac{1}{G_F} = \frac{N_c m_t^2}{4\pi^2 \sqrt{2}} \ln \frac{\Lambda_L^2}{m_t^2}.$$

Using our estimated value

$$\Lambda_L = 10^{14} \text{GeV}$$

and the measured value of the Fermi constant, we find the top mass

In particular, such relations give

$$\frac{1}{G_F} = \frac{N_c m_t^2}{4\pi^2 \sqrt{2}} \ln \frac{\Lambda_L^2}{m_t^2}.$$

Using our estimated value

$$\Lambda_L = 10^{14} \text{GeV}$$

and the measured value of the Fermi constant, we find the top mass

$$m_t = 171.6 \text{GeV}$$

In particular, such relations give

$$\frac{1}{G_F} = \frac{N_c m_t^2}{4\pi^2 \sqrt{2}} \ln \frac{\Lambda_L^2}{m_t^2}.$$

Using our estimated value

$$\Lambda_L = 10^{14} \text{GeV}$$

and the measured value of the Fermi constant, we find the top mass

$$m_t = 171.6 \text{GeV}$$

Moreover,

$$\rho \equiv \frac{\tilde{g}^2 m_W^2}{g^2 m_Z^2} = \frac{f_W}{f_Z} = 2 \frac{m_t^2 f'_{tb} + m_b^2 f'_{bt}}{m_t^2 f_{tt} + m_b^2 f_{bb}} \sim 1$$

for $\Lambda_L \gg m_t \gg m_b$

Conclusions

Conclusions

A weighted power-counting criterion can be used to renormalize Lorentz violating theories that contain **higher space derivatives**, but **no higher time derivatives**

Conclusions

A weighted power-counting criterion can be used to renormalize Lorentz violating theories that contain **higher space derivatives**, but **no higher time derivatives**

Renormalizable theories can contain two scalar-two fermion vertices and four fermion vertices

Conclusions

A weighted power-counting criterion can be used to renormalize Lorentz violating theories that contain **higher space derivatives**, but **no higher time derivatives**

Renormalizable theories can contain two scalar-two fermion vertices and four fermion vertices

We can construct an extended Standard Model that **gives masses to left neutrinos** without introducing right neutrinos or other extra fields. We can also describe **proton decay**

Conclusions

A weighted power-counting criterion can be used to renormalize Lorentz violating theories that contain **higher space derivatives**, but **no higher time derivatives**

Renormalizable theories can contain two scalar-two fermion vertices and four fermion vertices

We can construct an extended Standard Model that **gives masses to left neutrinos** without introducing right neutrinos or other extra fields. We can also describe **proton decay**

At very high energies gauge and Higgs interactions become negligible, since they are super-renormalizable, so the model becomes a four fermion model in two weighted dimensions.

Conclusions

A weighted power-counting criterion can be used to renormalize Lorentz violating theories that contain **higher space derivatives**, but **no higher time derivatives**

Renormalizable theories can contain two scalar-two fermion vertices and four fermion vertices

We can construct an extended Standard Model that **gives masses to left neutrinos** without introducing right neutrinos or other extra fields. We can also describe **proton decay**

At very high energies gauge and Higgs interactions become negligible, since they are super-renormalizable, so the model becomes a four fermion model in two weighted dimensions.

It is thus natural to ask ourselves if the scalarless variant of the model is able to reproduce all known low-energy physics without the ambiguities of usual Nambu—Jona-Lasinio non-renormalizable approaches.

Conclusions

A weighted power-counting criterion can be used to renormalize Lorentz violating theories that contain **higher space derivatives**, but **no higher time derivatives**

Renormalizable theories can contain two scalar-two fermion vertices and four fermion vertices

We can construct an extended Standard Model that **gives masses to left neutrinos** without introducing right neutrinos or other extra fields. We can also describe **proton decay**

At very high energies gauge and Higgs interactions become negligible, since they are super-renormalizable, so the model becomes a four fermion model in two weighted dimensions.

It is thus natural to ask ourselves if the scalarless variant of the model is able to reproduce all known low-energy physics without the ambiguities of usual Nambu—Jona-Lasinio non-renormalizable approaches.

We have shown, in the leading order of the large N_c limit and with gauge interactions switched off, that the effective potential admits a Lorentz invariant (local) minimum, that gives masses to fermion and gauge bosons, and produces composite Higgs bosons.

We shown that the scalarless model is unambiguous and predicts relations among the parameters of the Standard Model. Our approximation has a good 50% of error, but one prediction, the relation between the top mass and the Fermi constant turns out to be in astonishing agreement with experiment.

We shown that the scalarless model is unambiguous and predicts relations among the parameters of the Standard Model. Our approximation has a good 50% of error, but one prediction, the relation between the top mass and the Fermi constant turns out to be in astonishing agreement with experiment.

High-energy Lorentz violations could allow us to define the **ultraviolet limit** of quantum gravity. Hopefully suitable mechanisms could make the violations **undetectable** even in principle

We shown that the scalarless model is unambiguous and predicts relations among the parameters of the Standard Model. Our approximation has a good 50% of error, but one prediction, the relation between the top mass and the Fermi constant turns out to be in astonishing agreement with experiment.

High-energy Lorentz violations could allow us to define the **ultraviolet limit** of quantum gravity. Hopefully suitable mechanisms could make the violations **undetectable** even in principle

Observe that $\bar{d} = 2$ is the weighted dimension where Lorentz violating gravity becomes strictly renormalizable. The real challenge is to break the local Lorentz symmetry without destroying unitarity.

We shown that the scalarless model is unambiguous and predicts relations among the parameters of the Standard Model. Our approximation has a good 50% of error, but one prediction, the relation between the top mass and the Fermi constant turns out to be in astonishing agreement with experiment.

High-energy Lorentz violations could allow us to define the **ultraviolet limit** of quantum gravity. Hopefully suitable mechanisms could make the violations **undetectable** even in principle

Observe that $\bar{d} = 2$ is the weighted dimension where Lorentz violating gravity becomes strictly renormalizable. The real challenge is to break the local Lorentz symmetry without destroying unitarity.

If this were shown to be not possible, we would have a good reason to think that Lorentz invariance (and therefore CPT) must be exact at arbitrarily high energies.